

**СРПСКО КРИСТАЛОГРАФСКО ДРУШТВО
SERBIAN CRYSTALLOGRAPHIC SOCIETY**

**СРПСКО
КРИСТАЛОГРАФСКО ДРУШТВО
XI КОНФЕРЕНЦИЈА
СРПСКОГ КРИСТАЛОГРАФСКОГ ДРУШТВА
Изводи радова**

**XI CONFERENCE OF THE
SERBIAN CRYSTALLOGRAPHIC SOCIETY
Abstracts**

**SERBIAN CRYSTALLOGRAPHIC
SOCIETY**

ОПЛЕНАЦ - OPLENAC

2003

ISSN 0354-5741

СРПСКО КРИСТАЛОГРАФСКО ДРУШТВО
SERBIAN CRYSTALLOGRAPHIC SOCIETY

ХІ КОНФЕРЕНЦИЈА
СРПСКОГ КРИСТАЛОГРАФСКОГ ДРУШТВА
Изводи радова

XI CONFERENCE OF THE
SERBIAN CRYSTALLOGRAPHIC SOCIETY
Abstracts

ОПЛЕНАЦ - OPLENAC
2003

ХІ КОНФЕРЕНЦИЈА СРПСКОГ КРИСТАЛОГРАФСКОГ ДРУШТВА
Изводи радова

XI CONFERENCE OF THE SERBIAN CRYSTALLOGRAPHIC SOCIETY

Abstracts

Издавач – Publisher

- Српско Кристалографско Друштво

Ђушина 7, 11000 Београд, Србија и Црна Гора, тел/факс: 635 – 217

- Serbian Crystallographic Society

Đušina 7, 11000 Belgrade, Serbia and Montenegro, phone/fax: +381 – 11 – 635 – 217

За издавача – For the publisher

Љиљана Каравановић – Ljiljana Karanović

Технички уредник – Technical editor

Љиљана Каравановић – Ljiljana Karanović

Предраг Вулић – Predrag Vujić

Издавање ове публикације омогућено је финансијском помоћу Министарства за науку, технологије и развој Републике Србије

This publication is financially supported by Ministry of Science, Technology and Development of Republic of Serbia

© Српско Кристалографско Друштво – Serbian Crystallographic Society

ISSN 0354-5741

Штампа – Printing

СИГНАТУРЕ, Београд

SIGNATURE, Belgrade

Тираж – Copies: 100

Београд – Belgrade

2003

**XI КОНФЕРЕНЦИЈА
СРПСКОГ КРИСТАЛОГРАФСКОГ ДРУШТВА
Опленац, 25 – 27. септембар 2003.**

**XI CONFERENCE OF THE
SERBIAN CRYSTALLOGRAPHIC SOCIETY
Oplenac, 25 – 27 September 2003**

НАУЧНИ ОДБОР/SCIENTIFIC COMMITTEE:

- S. Stanković, PMF, Novi Sad
- A. Kapor, PMF, Novi Sad
- D. Poleti, TMF, Beograd
- B. Antić, INN „Vinča”, Beograd

ОРГАНИЗАЦИОНИ ОДБОР/ORGANIZING COMMITTEE:

- Lj. Karanović, RGF, Beograd
- G. Bogdanović, INN „Vinča”, Beograd
- A. Kremenović, RGF, Beograd
- P. Vulić, RGF, Beograd

CONTENT-САДРЖАЈ

ORAL PRESENTATIONS - САОПШТЕЊА

1. M. Milčić, Z. Tomić, S. Zarić STRONG METAL LIGAND AROMATIC CATION – π INTERACTIONS IN TETRAPHENYLBORATE SALTS.....	2
M. Milčić, Z. Tomić, S. Zarić JAKE METAL LIGAND AROMATSKE KATJON – π -INTERAKCIJE U SOLIMA SA TETRAFENILBORATNIM ANJONOM	3
2. N. Bošnjaković-Pavlović, I. Correia, I. Tomaz, N. Bouhmaida, F. Avecilla, A. Spasojević-de Biré, U. B. Mioč, J. Pessoa, N. E. Ghermani STRUCTURE AND ELECTROSTATIC PROPERTIES OF A DECAVANADATE-CYTOSINE COMPLEX.....	4
N. Bošnjaković-Pavlović, I. Correia, I. Tomaz, N. Bouhmaida, F. Avecilla, A. Spasojević-de Biré, U. B. Mioč, J. Pessoa, N. E. Ghermani STRUKTURA I ELEKTROSTATIČKE OSOBINE KOMPLEKSA DECAVANADIJUM-CITOZIN	5
3. A. Bacchi, G. Pelizzi, G. Jakovljević, N. Filipović, M. Rajković, K. Andelković SYNTHESIS AND CRYSTAL STRUCTURE OF TRIS[(1-PYRIDIN-2-YL- ETHYLIDENE)HYDRAZINE]IRON(II) PERCHLORATE	6
A. Bacchi, G. Pelizzi, G. Jakovljević, N. Filipović, M. Rajković, K. Andelković SINTEZA I KRISTALNA STRUKTURA TRIS[(1-PIRIDIN-2-IL-ETILIDEN)HIDRAZIN]GVOŽĐE(II)- PERHLORATA.....	7
4. V. B. Medaković, Lj. Karanović, D. Poleti, V. M. Leovac, S. D. Zarić ORIENTATION OF COORDINATED PYRIDINE IN CRYSTAL STRUCTURE OF COBALT(III) COMPLEX WITH SALICYLALDEHYDE S-METHYLISOTHOSEMICARBAZONE	8
V. B. Medaković, Lj. Karanović, D. Poleti, V. M. Leovac, S. D. Zarić ORIJENTACIJA KOORDINOVANOG PIRIDINA U KRISTALNOJ STRUKTURI KOMPLEKSA KOBALTA(III) SA S-METILIZOTIOSEMIKARBAZONOM SALICILALDEHIDA.....	9
5. J. Rogan, D. Poleti, Lj. Karanović A BINUCLEAR COBALT(II) COMPLEX CONTAINING PYROMELLITATE ION AND 1,10- PHENANTHROLINE.....	10
J. Rogan, D. Poleti, Lj. Karanović BINUKLEARNI KOBALT(II) KOMPLEKS SA PIROMELITAT JONOM I 1,10-FENANTROLINOM	11
6. J. Rogan, D. Poleti, Lj. Karanović CRYSTAL STRUCTURE OF A NEW NICKEL(II) COMPLEX CONTAINING PYROMELLITATE ION AND 2,2'-DIPYRIDYLAMINE.....	12
J. Rogan, D. Poleti, Lj. Karanović KRISTALNA STRUKTURA NOVOG NIKAL(II) KOMPLEKSA SA PIROMELITAT JONOM I 2,2'-DIPIRIDIILAMINOM	13
7. S. B. Novaković, G. A. Bogdanović, V. M. Leovac CRYSTAL STRUCTURE OF THE Ni(II) COMPLEX WITH S-METHYLISOTHOSEMICARBAZIDE AND TEREPHTHALATE.....	14
S. B. Novaković, G. A. Bogdanović, V. M. Leovac KRISTALNA STRUKTURA KOMPLEKSA Ni(II) SA S-METILIZOTIOSEMIKARBAZIDOM I TEREFTALATOM	15
8. Z. D. Tomić, S. B. Novaković, S. D. Zarić ON THE FACTORS INFLUENCING THE INTERMOLECULAR CONTACTS BETWEEN THE COPPER(II) AND PHENYL RING IN THE SQUARE-PLANAR Cu(II) COMPLEXES	16
Z. D. Tomić, S. B. Novaković, S. D. Zarić FAKTOVI KOJI UTIČU NA INTERMOLEKULSKE KONTAKTE IZMEDJU BAKRA(II) I FENILNOG PRSTENA KOD KVADRATNO-PLANARNIH Cu(II) KOMPLEKSA	17

9. S. R. Trifunović, G. N. Kaluderović, F. W. Heinemann, T. J. Sabo	
CRYSTAL STRUCTURE OF POTASSIUM BIS(MALONATO)CUPRATE(II) DIHYDRATE.....	18
S. R. Trifunović, G. N. Kaluderović, F. W. Heinemann, T. J. Sabo	
KRISTALNA STRUKTURA KALIJUM-BIS(MALONATO)KUPRAT(II) DIHIDRATA.....	19
10. G. N. Kaluderović, D. Poleti, Lj. Karanović, T. J. Sabo	
CRYSTAL STRUCTURE OF <i>TRANS</i> -DIBROMO(ETHYLENEDIAMINE- <i>N,N'</i> -DI-3-PROPIONATO)PLATINUM(IV) MONOHYDRATE.....	20
G. N. Kaluderović, D. Poleti, Lj. Karanović, T. J. Sabo	
KRISTALNA STRUKTURA <i>TRANS</i> -DIBROMO(ETILENDIAMIN- <i>N,N'</i> -DI-3-PROPIONATO)PLATINA(IV) MONOHIDRATA.....	21
11. T. Soldatović, F. W. Heinemann, Ž. D. Bugarčić	
SYNTHESIS AND CRUSTAL STRUCTURE OF [Pt(terpy)(cyst-S)][ClO ₄] ₂ ·0.5H ₂ O.....	22
T. Soldatović, F. W. Heinemann, Ž. D. Bugarčić	
SINTEZA I KRISTALNA STRUKTURA [Pt(terpy)(cyst-S)][ClO ₄] ₂ ·0.5H ₂ O KOMPLEKSA	23
12. B. Petrović, E. Zangrando, Ž. D. Bugarčić	
SYNTHESIS AND CRUSTAL STRUCTURE OF [Pd(terpy)(py)][ClO ₄] ₂	24
B. Petrović, E. Zangrando, Ž. D. Bugarčić	
SINTEZA I KRISTALNA STRUKTURA [Pd(terpy)(py)][ClO ₄] ₂ KOMPLEKSA.....	25
13. T. Mihajlović, H. Effenberger, E. Tillmanns	
BaMgAs ₂ O ₇ AND BaCoAs ₂ O ₇ – SYNTHESIS AND CRYSTAL STRUCTURES OF TWO NEW BARIUM DIARSENATES.....	26
T. Mihajlović, H. Effenberger, E. Tillmanns	
BaMgAs ₂ O ₇ I BaCoAs ₂ O ₇ – SINTEZA I KRISTALNA STRUKTURA DVA NOVA DIARSENATA BARIJUMA.....	27
14. G. Berti	
THE STATE OF ART AND EXPECTED DEVELOPMENTS ON THE CALIBRATION OF DIFFRACTOMETER.....	28
15. M. Mitrić, J. Blanuša, Z. Dohčević-Mitrović, N. Paunović, V. Spasojević	
STRUCTURAL PROPERTIES OF La _{1-x} Na _x MnO ₃ SOLID SOLUTIONS	32
M. Mitrić, J. Blanuša, Z. Dohčević-Mitrović, N. Paunović, V. Spasojević	
STRUKTURNYE OSOBINE ČVRSTIH RASTVORA La _{1-x} Na _x MnO ₃	33
16. A. Rosić, S. Radulović	
DETERMINATION OF MINERAL COMPOSITION OF URINARY CALCULI BY X-RAY POWDER DIFFRACTION ANALYSIS.....	34
A. Rosić, С. Радуловић	
ОДРЕЂИВАЊЕ МИНЕРАЛНОГ САСТАВА УРИНАРНОГ КАМЕЊА МЕТОДОМ РЕНДГЕНСКЕ ДИФРАКЦИЈЕ ПРАХА.....	35
17. A. Kremenović, B. Antić, A. S. Nikolić, M. Stoiljković	
STRUCTURE AND MICROSTRUCTURE OF CATION DEFICIT Zn-Mn FERRITES	36
A. Kremenović, B. Antić, A. S. Nikolić, M. Stoiljković	
STRUKTURA I MIKROSTRUKTURA KATJONSKI DEFICITARNIH Zn-Mn FERITA	37
18. P. Vulić, R. Dimitrijević, V. Dondur, S. Marković	
MICROSTRUCTURE PARAMETERS OF Na-NEPHELINES WITH DIFFERENT Si/Al RATIO	38
П. Вулић, Р. Димитријевић, В. Дондур, С. Марковић	
МИКРОСТРУКТУРНИ ПАРАМЕТРИ Na-НЕФЕЛИНА СА РАЗЛИЧИТИМ ОДНОСОМ Si/Al	39
19. M. Zdujić, Č. Jovalekić, Lj. Karanović, D. Poleti	
2Bi ₂ O ₃ ·3TiO ₂ SYSTEM: STRUCTURAL CHANGES INDUCED BY INTENSIVE MILLING.....	40
М. Здујић, Ч. Јовалекић, Љ. Карановић, Д. Полети	
2Bi ₂ O ₃ ·3TiO ₂ СИСТЕМ: СТРУКТУРНЕ ПРОМЕНЕ ИНДУКОВАНЕ ИНТЕНЗИВНИМ МЛЕВЕЊЕМ	41

20. A. Bordaš, S. Rakić, B. Antić, A. Kapor	
RADIAL DISTRIBUTION FUNCTION AND THE LOCAL STRUCTURE OF SOLID SOLUTIONS OF Li _{0.5} Fe _{2.5-x} La _x O ₄ TYPE.....	42
A. Bordaš, S. Rakić, B. Antić, A. Kapor	
RADIJALNA FUNKCIJA RASPODELE I LOKALNA STRUKTURA ČVRSTIH RASTVORA TIPA Li _{0.5} Fe _{2.5-x} La _x O ₄	43
21. A. Kapor, D. Zobel, M. Strümpel, Lj. Torović, M. Popsavin	
NEKOVALENTE INTERAKCIJE U KRISTALNOJ STRUKTURI E- I Z- IZOMERA NEZASIĆENOG ESTRA C-GLUKOZIDA.....	44
A. Kapor, D. Zobel, M. Strümpel, Lj. Torović, M. Popsavin	
NON-COVALENT INTERACTION IN CRYSTAL STRUCTURE OF E- AND Z- ISOMERS OF UNSATURATED ESTERS OF C-GLYCOSIDES	45
22. G. A. Bogdanović, V. S. Jevtović, V. M. Leovac	
THREE-DIMENSIONAL HYDROGEN-BONDING NETWORK IN THE CRYSTAL STRUCTURE OF PYRIDOXAL SEMICARBAZONE CHLORIDE MONOHYDRATE	46
G. A. Bogdanović, V. S. Jevtović, V. M. Leovac	
TRODIMENZIJALNA MREŽA VODONIČNIH VEZA U KRISTALNOJ STRUKTURI PIRIDOKSAL- SEMIKARBAZON-HLORID-MONOHIRATA	47
23. D. Lazar, S. Stanković, O. Klisurić, S. Jovanović-Šanta, J. Petrović	
SYNTHESIS AND CRYSTAL STRUCTURE OF 3,16,17-TRIHYDROXY-16,17-SECOESTRA-1,3,5(10)- TRIENE	48
D. Lazar, S. Stanković, O. Klisurić, S. Jovanović-Šanta, J. Petrović	
SINTEZA I KRISTALNA STRUKTURA 3,16,17-TRIHDROKSI-16,17-SEKOESTRA-1,3,5,(10)-TRIENA	49
24. M. Garić, D. Ž. Obadović, S. Jovanović-Šanta, D. Lazar, A. Vajda	
CRYSTAL STRUCTURE AND THE INFLUENCE OF NEW 3,17-SUBSTITUED D-SECO-ESTRONE DERIVATIVES ONTO THE PHYSICAL CHARACTERISTICS OF THE BINARY MIXTURES OF THE CHOLESTERIC LIQUID CRYSTALS	50
M. Garić, D. Ž. Obadović, S. Jovanović-Šanta, D. Lazar, A. Vajda	
KRISTALNA STRUKTURA I UTICAJ NOVIH 3,17-SUPSTITUISANIH D-SEKO-ESTRONSKIH DERIVATA NA FIZIČKE KARAKTERISTIKE BINARNIH SMEŠA HOLESTERIČKIH TEČNIH KRISTALA	51
25. O. Klisurić, S. Stanković, D. Lazar, D. Zobel, K. Penov-Gaši, S. Stajanović	
SYNTHESIS AND CRYSTAL STRUCTURE INVESTIGATIONS OF TWO ANDROSTENE DERIVATIVES	52
O. Klisurić, S. Stanković, D. Lazar, D. Zobel, K. Penov-Gaši, S. Stajanović	
SINTEZA, STRUKTURNA I KRISTALOGRAFSKA ISTRAŽIVANJA DVA DERIVATA ANDROSTENA	53
26. O. Klisurić, S. Stanković, D. Lazar, D. Zobel, A. Wagner, K. Penov-Gaši, S. Stajanović	
SYNTHESIS AND CRYSTAL STRUCTURE OF (17R)-3,17-DIHYDROXY-16-CYANO-17-METHYL-16,17-SECOESTRA-1,3,5(10)-TRIENE (1) AND 3-BENZYLOXY-17A-METHYL-17-OXA-D-HOMOESTRA-1,3,5-TRIEN-16-ONE (2)	54
O. Klisurić, S. Stanković, D. Lazar, D. Zobel, A. Wagner, K. Penov-Gaši, S. Stajanović	
SINTEZA I KRISTALNA STRUKTURA (17R)-3,17-DIHDROKSI-16-CIJANO-17-METIL-16,17- SEKOESTRA-1,3,5(10)-TRIENA (1) I 3-BENZILOKSI-17A-METIL-17-OKSA-D-HOMO-ESTRA-1,3,5- TRIEN-16-ONA (2)	55

ORAL PRESENTATIONS

САОПШТЕЊА

STRONG METAL LIGAND AROMATIC CATION – π INTERACTIONS IN TETRAPHENYLBORATE SALTS

M. Milčić^a, Z. Tomic^b and S. Zarić^a

^aDepartment of Chemistry, University of Belgrade, Studentski trg 16, P.O. Box 158, 11001 Belgrade, Serbia and Montenegro; ^b"Vinča" Institute of Nuclear Sciences, Laboratory of Theoretical Physics and Condensed Matter Physics, 11001 Belgrade, P.O. Box 522, Serbia and Montenegro
e-mail: szaric@chem.bg.ac.yu

The cation – π interaction is considered to be an important non-covalent bonding force in a wide range of molecular systems. It was shown that this type of interactions can also exist when the cation is a transition metal complex [1]. Cation- π interactions between ligands coordinated to a metal cation and aromatic groups, metal ligand aromatic cation – π interactions (MLAC- π), have been found in crystal structures of metalloproteins from the Protein Data Bank [2] and as intramolecular interaction, in crystal structures of transition metal complexes from Cambridge Structural Database (CSD) [3].

Since in MLAC π interactions hydrogen atoms from ligands interact with π -systems, this interaction can be considered to be X-H... π bond. However, it is charge assisted bond since, by coordinating to the metal cation, ligands gain some positive charge. Here we report results of searching for intermolecular MLAC- π interactions between the hydrogen atom from a positively charged transition metal complex and phenyl groups from tetraphenylborate (TFB) anion. Since TFB is negatively charged it can be anticipated that this interactions can be relatively strong.

Screening the CSD provided over 200 structures in which an aromatic ring from TFB is close to a metal. We inspected these structures and selected 15 structures where the X... π c (π c - center of phenyl ring) distances were very short, and at the same time the X-H... π c angle was larger than 100°. Based on the number of phenyl groups from the TFB anion interacting with the same ligand from metal complex, there are two types of intermolecular MLAC- π interactions. In the first type (I) two phenyl groups from TFB anion interact with the same ligand. In the second type (II) one aromatic group from the TFB anion interact with a ligand from transition metal complex.

- [1] S. D. Zarić, *Chem. Phys. Lett.*, 311 (1999), 77.
- [2] S. D. Zarić, D. M. Popović, E. W. Knapp, *Chemistry*, 6 (2000), 3935.
- [3] M. K. Milčić, S. D. Zarić, *Eur.J.Inorg.Chem.*, (2001), 2143.

JAKE METAL LIGAND AROMATSKE KATJON – π -INTERAKCIJE U SOLIMA SA TETRAFENILBORATNIM ANJONOM

M. Milčić^a, Z. Tomić^b i S. Zarić^a

^aHemski fakultet, Univerzitet u Beogradu, Studentski trg 16, P.O. Box 158, 11001 Beograd, Srbija i Crna Gora; ^b"Vinča" Institut za nuklearne nauke, Laboratorija za teorijsku fiziku i fiziku čvrstog stanja, 11001 Beograd, P.O. Box 522, Srbija i Crna Gora
e-mail: szaric@chem.bg.ac.yu

Katjon – π -interakcije se smatraju veoma bitnim nekovalantnim vezivnim interakcijama u velikom broju molekulskih sistema. Pokazano je da te interakcije postoje i u sistemima u kojim je katjon kompleks prelaznog metala [1]. Katjon – π -interakcije između liganada koordinovanih za metalni katjon i aromatičnih grupa, metal ligand aromatske katjon- π interakcije (MLAC- π), su pronađene u kristalnim strukturama metaloproteina iz proteinske banke podataka (PDB) [2] i, kao intremolekulske interakcije, u kristalnim strukturama kompleksa prelaznih metala iz Kembridžske banke kristalografskih podataka (CSD) [3].

Pošto u ovom tipu interakcija vodonikov atom sa ligandom interaguje sa π -sistom ove interakcije se mogu posmatrati kao X-H... π veza. Ova veza nastaje zbog toga što koordinacijom za centralni metalni ion, deo pozitivnog naelektrisanja prelazi sa metalnog jona na ligand. Ovde su izneti rezultati pretraživanja kristalografske banke podataka za intermolekulske MLAC- π interakcije između vodonikovog atoma sa pozitivno naelektrisanog kompleksa prelaznih metala i fenil grupe sa tetrafenilboratnog (TFB) anjona. Pošto je TFB negativno naelektrisan može se predpostaviti da će ove interakcije biti veoma jake.

Pretraživanjem banke kristalografskih podataka dobijeno je preko 200 struktura u kojima je aromatična grupa sa TFB anjona blizu metalnog jona. Sve ove strukture smo pobliže ispitali i odabrali njih 15 gde su X... π c (π c-centar benzenovog prstena) rastojanja veoma kratka, a u isto vreme ugao X-H... π c je veći od 100° . Na osnovu broja fenil grupa sa TFB anjona koje interaguju sa istim ligandom sa metalnog kompleksa, postoje dva tipa intermolekulske MLAC- π interakcija. U prvom tipu (I) ovih interakcija dve fenil grupe sa TFB anjona interaguju sa istim ligandom sa kompleksa prelaznog metala. U drugom tipu (II) samo jedna aromatična grupa sa TFB anjona interaguje sa jednim ligandom sa kompleksa prelaznog metala.

- [1] S. D. Zarić, *Chem. Phys. Lett.*, 311 (1999), 77.
- [2] S. D. Zarić, D. M. Popović, E. W. Knapp, *Chemistry*, 6 (2000), 3935.
- [3] M. K. Milčić, S. D. Zarić, *Eur.J.Inorg.Chem.*, (2001), 2143.

STRUCTURE AND ELECTROSTATIC PROPERTIES OF A DECAVANADATE-CYTOSINE COMPLEX

N. Bošnjaković-Pavlović^a, I. Correia^b, I. Tomaz^b, N. Bouhmaida^{a,c}, F. Avecilla^d, A. Spasojević-de Biré^a, U.B. Mioć^e, J. Pessoa^b et N. E. Ghermani^{a,f}

^aEcole Centrale Paris, Laboratoire SPMS UMR CNRS 8580 1, Grande Voie des Vignes 92295 Châtenay-Malabry, France; ^bCentro de Química Estrutural, Instituto Superior Técnico, Universidade Técnica de Lisboa, Av. Rovisco Pais, P-1049-001 Lisboa, Portugal; ^cLaboratoire des Sciences des Matériaux, LSM, Université Cadi Ayyad, Faculté des Sciences Semlalia, Boulevard Prince Moulay Abdallah, BP 2390, 40000 Marrakech, Maroc; ^dDepartamento de Química Fundamental, Facultad de Ciencias, Universidad de Coruña, Campus da Zapateira s/n, 15071 A Coruña, Espagne; ^eFaculty of Physical Chemistry, University of Belgrade, P.O.Box 137, 11001 Belgrade, Serbia and Montenegro; ^fLaboratoire de Physique Pharmaceutique UMR CNRS 8612, Faculté de Pharmacie 5, Rue Jean-Baptiste Clément, 92296 Châtenay-Malabry cedex, France
e-mail: nada@spms.ecp.fr

The polyoxometalates (POM's) of general formula $H_{6-x}A_xM_{10}O_{28}\cdot nH_2O$, ($A = K^+$, Na^+ , $M = W, V$, $n = 0-30$) crystallize with a various number of water molecules. POM's primary structure is formed with polyanions while the secondary structure is formed with polyanions, cations and water molecules. POM's are studied for their potentially pharmaceutical applications (antiviral or antitumoral). These biological properties result from the POM's interactions with enzymes of cells at the nucleus membranes. The study of complex of POM's with amino acids or nucleic bases is therefore of a particular interest for a better understanding of their therapeutic impact.

Among the different synthesized POM's, we have focused on the polyoxovanadates (POV's). The vanadium has been chosen for its relative low toxicity in biological media compared for example to the tungstene. It has been established that, POV's are inhibitors or activators in the enzymatic reactions as kynase [1], phosphorilase [2] and ATP-ase of Ca^{2+} [2].

We are reporting the preliminary results of the electron density and the electrostatic properties of a decavanadate-cytosine complex of formula $Na_3V_{10}O_{28}(C_4N_3OH_5)_3(C_4N_3OH_6)_3\cdot 10H_2O$ (**1**). These properties are obtained from high resolution X-ray diffraction data (0.44 Å) at 210 K. **1** crystallizes in the $P\bar{1}$ space group, the crystallographic structure is also governed by interactions between the cytosine molecules which are protonated or not. The protonation-deprotonation process gives rise to additional diffraction peaks at temperature lower than 210 K. An instable $P\bar{1}$ to $P1$ phase transition was observed. Diffraction data have been refined using the Hansen-Coppens multipole model [3]. The electrostatic properties of the decavanadate-cytosine complex will be presented and discussed.

- [1] D.W. Boyd, K. Kustin, M. Niwai, *Biochim. Biophys. Acta*, 827 (1985), 472.
- [2] P. Csermely, A. Martonosi, G. C. Levy, A. J. Ejchart, *Biochem. J.*, 230 (1985), 807.
- [3] N. Hansen and P. Coppens, *Acta Crystallogr.*, A34 (1978), 909.

STRUKTURA I ELEKTROSTATIČKE OSOBINE KOMPLEKSA DEKAVANADIJUM-CITOZINA

N. Bošnjaković-Pavlović^a, I. Correia^b, I. Tomaz^b, N. Bouhmaida^{a,c}, F. Avecilla^d, A. Spasojević-de Biré^a, U.B. Mioč^e, J. Pessoa^b, N. E. Ghermani^{a,f}

^aEcole Centrale Paris, Laboratoire SPMS UMR CNRS 8580 1, Grande Voie des Vignes 92295 Châtenay-Malabry, France; ^bCentro de Química Estrutural, Instituto Superior Técnico, Universidade Técnica de Lisboa, Av. Rovisco Pais, P-1049-001 Lisboa, Portugal; ^cLaboratoire des Sciences des Matériaux, LSM, Université Cadi Ayyad, Faculté des Sciences Semlalia, Boulevard Prince Moulay Abdallah, BP 2390, 40000 Marrakech, Maroc; ^dDepartamento de Química Fundamental, Facultad de Ciencias, Universidad de Coruña, Campus da Zapateira s/n, 15071 A Coruña, Espagne; ^eFakultet za Fizičku hemiju, Univerzitet u Beogradu, P.O.Box 137, 11001 Beograd, Srbija i Crna Gora; ^fLaboratoire de Physique Pharmaceutique UMR CNRS 8612, Faculté de Pharmacie 5, Rue Jean-Baptiste Clément, 92296 Châtenay-Malabry cedex, France
e-mail: nada@spms.ecp.fr, nada@ffh.bg.ac.yu

Poliosmetalati (POM) su jedinjenja opšte formule $H_{6-x}A_xM_{10}O_{28}\cdot nH_2O$, ($A = K^+$, Na^+ , $M = W$, V , $n = 0-30$) koja kristališu sa različitim brojem molekula vode. Primarna struktura POM-a je sastavljena od polianjona dok je sekundarna struktura sastavljena od polianjona, katjona i molekula vode. POM-i su ispitivani zbog njihove potencijalne primene u farmaciji (anti-tumorna i anti-viralna aktivnost). Njihove biološke osobine su rezultat interakcije POM-a sa enzimima u membranama ćelijskog jezgra. Ispitivanje kompleksa POM-a sa amino kiselinama ili nukleinskim bazama je od ključnog interesa za bolje razumevanje njihovih terapeutskih dejstava.

Između različito sintetisanih POM-a, mi smo fokusirali naša ispitivanja na polioksovanadatima (POV). Vanadijum smo izabrali zbog relativno manje toksičnosti u biološkom medijumu u poređenu sa volframom. POV su inhibitori ili aktivatori u reakcijama enzima kao što su kinaze [1], fosforilaze [2] i ATP-aze [2].

Mi ćemo predstaviti preliminarne rezultate određivanja elektronske gustine kao i elektrostatičke osobine kompleksa dekavanadijum-citozina, formule $Na_3V_{10}O_{28}(C_4N_3OH_5)_3(C_4N_3OH_6)_3\cdot 10H_2O$ (**1**). Ovi rezultati su dobijeni na osnovu merenja visoke rezolucije (0.44 Å) intenziteta difrakcionih maksimuma sa monokristala na temperaturi od 210 K. Jedinjenje **1** kristalise u $P\bar{1}$ prostornoj grupi i njegova kristalna struktura je pod uticajem interakcija koje međusobno grade molekuli citozina. Na temperaturama nižim od 210 K je opažena pojava dodatnih difrakcionih maksimuma koji su posledica mogućnosti protonovanja i deprotonovanja molekula citozina. Opažen je nestabilan fazni prelaz iz prostorne grupe $P\bar{1}$ u $P1$. Difrakcionni podaci su utačnjeni uz korišćenje Hansen-Coppens multipol modela [1]. Elektrostatičke osobine kompleksa dekavanadijum-citozina će biti prezentirane i diskutovane.

- [1] D.W. Boyd, K. Kustin, M. Niwai, *Biochim. Biophys. Acta*, 827 (1985), 472.
- [2] P. Csermely, A. Martonosi, G. C. Levy, A. J. Ejchart, *Biochem. J.*, 230 (1985), 807.
- [3] N. Hansen and P. Coppens, *Acta Crystallogr.*, A34 (1978), 909.

SYNTHESIS AND CRYSTAL STRUCTURE OF TRIS[(1-PYRIDIN-2-YL-ETHYLIDENE)HYDRAZINE]IRON(II) PERCHLORATE


A. Bacchi^a, G. Pelizzi^a, G. Jakovljević^b, N. Filipović^c, M. Rajković^c and K. Andelković^b

^a Dipartimento di Chimica Generale ed Inorganica, Chimica Analitica, Chimica Fisica, University of Parma, Parco Area delle Scienze 17A, I-43100 Parma, Italy; ^b Faculty of Chemistry, University of Belgrade, P. O. Box 158, 11001 Belgrade, Serbia and Montenegro; ^c Faculty of Agriculture, University of Belgrade, Nemanjina 6, 11081 Belgrade-Zemun, Serbia and Montenegro
e-mail: kka@chem.bg.ac.yu

The suspension of 0.42g (0.9 mmol) $\text{Fe}(\text{ClO}_4)_3 \cdot 6\text{H}_2\text{O}$ dissolved in 25 cm³ water, 0.16g (1.35 mmol) dihydrazide oxalic acid and 0.33g (2.7 mmol) 2-acetylpyridine (molar ratio 2:3:6) together with 3 drops of HClO_4 ($c = 0.1 \text{ mol/dm}^3$) was exposed to heating for 30 minutes, and dark-red microcrystalline product was obtained in a yield of 45%.

During the coordination reaction, oxidative degradation of -CO-CO- side chain fragment and reduction Fe(III) to Fe(II) occurred. Besides that, a molecule of carbon-dioxide was obtained in the reaction. The structure of Fe(II) complex is given in a Figure 1.

Crystal data: $\text{C}_{21}\text{H}_{27}\text{Cl}_2\text{FeN}_9\text{O}_8$, $M_W = 660.27$, radiation MoK α , $\lambda = 0.71073 \text{ \AA}$, monoclinic, $C2/c$, $a = 36.023(5)$, $b = 14.349(3)$, $c = 10.612(2) \text{ \AA}$, $\beta = 96.36(2)^\circ$, $V = 5451.5(17) \text{ \AA}^3$, $Z = 8$, Density (calculated) = 1.609 Mg/m^3 , Reflections collected = 5735, Independent reflections = 3129 [$R(\text{int}) = 0.1192$], Data / restraints / parameters = 3129 / 5 / 393, Final R indices [$I > 2\sigma(I)$]: $R_1 = 0.0739$, $wR_2 = 0.1658$.



SINTEZA I KRISTALNA STRUKTURA TRIS[(1-PIRIDIN-2-IL-ETILIDEN)HIDRAZIN] GVOŽĐE(II) – PERHLORATA

A. Bacchi^a, G. Pelizzi^a, G. Jakovljević^b, N. Filipović^c, M. Rajković^c i K. Andelković^b


^a Dipartimento di Chimica Generale ed Inorganica, Chimica Analitica, Chimica Fisica, University of Parma, Parco Area delle Scienze 17A, I-43100 Parma, Italy; ^b Hemski fakultet, Univerzitet u Beogradu, P.O. Box 158, 11001 Beograd, Srbija i Crna Gora

^c Poljoprivredni fakultet, Universitet u Beogradu, Nemanjina 6, 11081 Beograd-Zemun, Srbija i Crna Gora
e-mail: kka@chem.bg.ac.yu

Polazeći od 0,42g (0,9 mmol) $\text{Fe}(\text{ClO}_4)_3 \cdot 6\text{H}_2\text{O}$ rastvorenog u 25 ml vode, 0,16g (1,35 mmol) dihidrazida oksalne kiseline i 0,33g (2,7 mmol) 2-acetylpiridina (molski odnos 2:3:6) uz dodatak 2-3 kapi HClO_4 ($c = 0,1 \text{ mol/dm}^3$) pri zagrevanju na 65°C oko 30 minuta dobijen je tamnocrveni monokristalni proizvod u prinosu od 45 %.

U toku reakcije koordinovanja došlo je do oksidacione degradacije fragmenta bočnog lanca (-CO-CO-) i redukcije gvoždja(III) do gvoždja(II). Pri tome se iz sistema izdvadio ugljen dioksid. Dobijeni kompleks gvoždja(II) ima strukturu prikazanu na slici 1.

Kristalografski podaci: $\text{C}_{21}\text{H}_{27}\text{Cl}_2\text{FeN}_9\text{O}_8$, $M_W = 660.27$, zračenje MoK α , $\lambda = 0,71073 \text{ \AA}$, monoklinični, C2/c, $a = 36,023(5)$, $b = 14,349(3)$, $c = 10,612(2) \text{ \AA}$, $\beta = 96,36(2)^\circ$, $V = 5451,5(17) \text{ \AA}^3$, $Z = 8$, gustina (izračunata) = $1,609 \text{ Mg/m}^3$, prikupljene refleksije: 5735, nezavisne refleksije: 3129 [R(int) = 0,1192], podaci / ograničenja / parametri = 3129 / 5 / 393, konačni R faktor [$I > 2\sigma(I)$]: R1 = 0,0739, wR2 = 0,1658.



ORIENTATION OF COORDINATED PYRIDINE IN CRYSTAL STRUCTURE OF COBALT(III) COMPLEX WITH SALICYLALDEHYDE S-METHYLISOTHOSEMICARBAZONE

V. B. Medaković^a, Lj. Karanović^b, D. Poletić^c, V. M. Leovac^d, S. D. Zarić^a

^a Faculty of Chemistry, University of Belgrade, P. O. Box 158, 11001 Belgrade; ^b Faculty of Mining and Geology, Dušina 7, Belgrade; ^c Faculty of Technology and Metallurgy, Karnegijeva 4, Belgrade; ^d Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 3, 21000 Novi Sad;
e-mail: szaric@chem.bg.ac.yu

Orientation of axially coordinated ligand can be very important in biomolecules. The orientation of histidines, axially ligated to heme, are considered to have influence on the function of heme cofactors in proteins. The orientation of histidines was studied by analyzing crystal structures of heme proteins [1] and by quantum chemical calculations [2]. It was shown that orientation of histidines is determined by interactions with side groups on porphyrin.

The orientations of the pyridines in $[\text{Co}(\text{L})(\text{py})_3]\text{I}_3$ (H_2L = salicylaldehyde S-methylisothiosemicarbazone, py = pyridine) were studied by quantum chemical calculations [3]. Orientations of pyridines are caused by mutual influences of pyridines.

In this work we present the crystal structure of $[\text{Co}(\text{L})(\text{bpy})(\text{py})]$ (bpy = bipyridine) where coordination around Co(III) is distorted octahedral. Salicylaldehyde S-methylisothiosemicarbazone ligand is ONN-tridentate coordinated. Orientation of axially coordinated pyridine is different than in $[\text{Co}(\text{L})(\text{py})_3]\text{I}_3$ complex.

Crystallographic data: $\text{C}_{24}\text{H}_{24}\text{CoI}_3\text{N}_6\text{OS}$, triclinic crystal system, space group P-1, $a = 9.990(3)$, $b = 10.063(2)$, $c = 14.739(3)$ Å, $\alpha = 96.063(6)$, $\beta = 105.217(5)$, $\gamma = 90.523(4)$ °, $V = 1420.7(6)$ Å³, $Z = 2$, $F(000) = 840$, $D_c = 2.067$ Mg/m³, MoKα - radiation, $\mu(\text{MoK}\alpha) = 3.966$ mm⁻¹.

- [1] S. D. Zarić, D. Popović, E. W. Knapp, *Biochemistry* 40 (2001), 7914-7928.
- [2] V. B. Medaković, S. D. Zarić, *Inorg. Chim. Acta*, 349 (2003), 1-5.
- [3] G. A. Bogdanović, V. B. Medaković, Lj. S. Vojinović, V. I. Češljević, V. M. Leovac, S. D. Zarić, *Polyhedron*, 20 (2001), 2231-2240.

ORIJENTACIJA KOORDINOVANOG PIRIDINA U KRISTALNOJ STRUKTURI KOMPLEKSA KOBALTA(III) SA S-METILIZOTIOSEMIKARBAZONOM SALICILALDEHIDA

V. B. Medaković^a, Lj. Karanović^b, D. Poleti^c, V. M. Leovac^d, S. D. Zarić^a

^a Hemijski fakultet, Univerzitet u Beogradu, p. pr. 158, 11001 Belgrade; ^b Rudarsko-geološki fakultet, Đušina 7, Beograd; ^c Tehnološko-metalurški fakultet, Carnegiea 4, Beograd; ^d Prirodno-matematički fakultet, Univerzitet u Novom Sadu, Trg Dositeja Obradovića 3, 21000 Novi Sad;

Orijentacija aksijalno koordinovanog liganda ima veoma važnu ulogu u biomolekulima. Orijentacija histidina, aksijalno vezanog za hem, jako utiče na funkciju kofaktora hema u proteinima. Orijentacija histidina proučavana je analizom kristalnih struktura proteina hema [1] i kvantno hemijskim proračunima [2]. Pokazano je da je orijentacija histidina određena interakcijama sa bočnim grupama iz porfirina.

Faktori koji utiču na orijentaciju koordinovanih piridina u $[\text{Co}(\text{L})(\text{py})_3]\text{I}_3$ ($\text{H}_2\text{L} = \text{S-metilizotiosemikarbazon salicilaldehida, py = piridin}$) proučavani su kvantno hemijskim metodama [3]. Orijentacije piridina uzrokovane su interakcijama između aksijalnih i ekvatorijalnog piridina.

U ovom radu predstavljena je kristalna struktura $[\text{Co}(\text{L})(\text{bpy})(\text{py})]\text{I}_3$, ($\text{bpy} = \text{bipiridin}$) gde koordinacija oko Co(III) ima geometriju iskrivljenog oktaedra. S-metilizotiosemikarbazon salicilaldehida koordinovan je kao tridentatni ligand. Orijentacija aksijalno koordinovanog piridina razlikuje se od orijentacije u kompleksu $[\text{Co}(\text{L})(\text{py})_3]\text{I}_3$.

Kristalografski podaci: $\text{C}_{24}\text{H}_{24}\text{CoI}_3\text{N}_6\text{OS}$, triklinični kristalni sistem, prostorna grupa P-1, $a = 9,990(3)$, $b = 10,063(2)$, $c = 14,739(3)$ Å, $\alpha = 96,063(6)$, $\beta = 105,217(5)$, $\gamma = 90,523(4)$ °, $V = 1420,7(6)$ Å³, $Z = 2$, $F(000) = 840$, $D_c = 2,067$ Mg/m³, MoKα - zračenje, $\mu(\text{MoK}\alpha) = 3,966$ mm⁻¹.

- [1] S. D. Zarić, D. Popović, E. W. Knapp, *Biochemistry* 40 (2001), 7914-7928.
- [2] V. B. Medaković, S. D. Zarić, *Inorg. Chim. Acta*, 349 (2003), 1-5.
- [3] G. A. Bogdanović, V. B. Medaković, Lj. S. Vojinović, V. I. Češljević, V. M. Leovac, S. D. Zarić, *Polyhedron*, 20 (2001), 2231-2240.

A BINUCLEAR COBALT(II) COMPLEX CONTAINING PYROMELLITATE ION AND 1,10-PHENANTHROLINE

J. Rogan^a, **D. Poleti**^a, **Lj. Karanović**^b

^a Faculty of Technology and Metallurgy, Karnegijeva 4, Belgrade, Serbia and Montenegro,

^b Faculty of Mining and Geology, Dušina 7, Belgrade, Serbia and Montenegro

e-mail: rogan@elab.tmf.bg.ac.yu

Described compound presents continuation of our studies on ternary transition metal complexes containing anions of benzenopolycarboxylic acids and some aromatic diamines, as 1,10-phenanthroline (phen) [1]. Due to the presence of four carboxylate groups, the tetraanion of 1,2,4,5-benzenetetracarboxylic (pyromellitic or pyr) acid can coordinate in numerous ways, acting as a bis monodentat [1] or polydentate [2,3] ligand having a bridging, chelating or combined function. In the polymeric complex $\{[Cu_2(pyr)(phen)_2] \cdot (H_2O)\}_n$, which has similar empirical formula to the title complex, pyr acts as tetradeятate ligand and four COO groups are coordinated to four Cu atoms forming a double chain structure [3].

On the contrary, in our binuclear complex $[Co_2(pyr)(phen)_2(H_2O)_6] \cdot 2H_2O$, pyr is coordinated as bis-monodentate ligand. In addition, two *ortho*-COO groups are uncoordinated. The centre of symmetry is coinciding with the centre of aromatic ring. Co(II) environment consists of two N atoms (from phen) and four O atoms (three from H_2O molecules and one from COO group). The coordination polyhedron can be described as deformed octahedron with bond distances between 2.0550(19) and 2.1893(17) Å, and angles in the range 83.06(7) – 99.52(7) °.

Three coordinated H_2O molecules are the double donors in H bond network. One of them is the acceptor, too. The lattice H_2O molecule is an acceptor (toward O atom from one coordinated H_2O molecule) and donor (toward one O atom from uncoordinated COO group). There are four intramolecular H bonds, 2.588(3) and 2.714(2) Å, between coordinated H_2O and O atoms from COO groups.

Crystal data: $[Co_2(pyr)(phen)_2(H_2O)_6] \cdot 2H_2O$, $C_{34}H_{34}Co_2N_4O_{16}$, $M_r = 872.51$, triclinic, space group $P\bar{1}$, $a = 7.410(5)$, $b = 11.067(5)$, $c = 12.219(5)$ Å, $\alpha = 66.638(5)$, $\beta = 76.402(5)$, $\gamma = 70.855(5)$ °, $V = 862.61(8)$ Å³, $Z = 1$, $F(000) = 448$, $\rho_x = 1.680$ g cm⁻³, $\mu(Mo K\alpha) = 1.047$ mm⁻¹, 4251 independent reflections, $R_1 = 3.29$ % for 3224 observed reflections with $I > 2\sigma(I)$, $R_{w2} = 7.17$ % (refinement on F^2) for all reflections and 285 refined parameters, with $(\Delta/\sigma)_{max} = 0.003$, $\Delta\rho_{max} = 0.323$ and $\Delta\rho_{min} = -0.286$ e Å⁻³.

- [1] D. Poleti, D. R. Stojaković, B. V. Prelesnik, R. M. Herak, *Acta Cryst.*, **C44** (1988), 242–245.
- [2] R.-K. Chiang, N.-T. Chuang, Ch.-Sh. Wur, M.-F. Chong, Ch.-R. Lin, *J. Solid State Chem.*, **166** (2002), 158–163.
- [3] Q. Shi, R. Cao, D.-F. Sun, M.-Ch. Hong, Y.-C. Liang, *Polyhedron*, **20** (2001), 3287–3293.

BINUKLEARNI KOBALT(II) KOMPLEKS SA PIROMELITAT JONOM I 1,10-FENANTROLINOM

J. Rogan^a, D. Poleti^a, Lj. Karanović^b

^a Tehnološko-metalurški fakultet, Karnegijeva 4, Beograd, Srbija i Crna Gora, ^b Rudarsko-geološki fakultet, Đušina 7, Beograd, Srbija i Crna Gora

e-mail: rogan@elab.tmf.bg.ac.yu

Opisano jedinjenje predstavlja nastavak naših istraživanja ternernih kompleksa prelaznih metala sa anjonima benzenpolikarboksilnih kiselina i nekih aromatičnih diammina, kao što je 1,10-fenatrolin (phen) [1]. Zbog prisustva četiri COO grupe, tetraanjon 1,2,4,5-benzentetrakarboksilne (piromelitne, pyr) kiseline može da se koordinira na brojne načine, ponašajući se kao bis monodentatni [1] ili polidentatni [2,3] ligand sa mostovnom, helatnom ili kombinovanom ulogom. U kompleksu $\{[Cu_2(pyr)(phen)_2] \cdot (H_2O)\}_n$, koji ima sličnu empirijsku formulu kao naše jedinjenje, pyr se ponaša kao tetradentatni ligand i četiri COO grupe su vezane za četiri Cu atoma gradeći dvostruke lance [3].

Suprotno ovome, u našem binuklearnom kompleksu, $[Co_2(pyr)(phen)_2(H_2O)_6] \cdot 2H_2O$, pyr je koordiniran kao bis-monodentatni ligand. Dodatno, dve orto-COO grupe su nekoordinirane. Centar simetrije se poklapa sa centrom aromatičnog prstena. Okruženje oko Co(II) čine dva N atoma (iz phen) i četiri O atoma (tri iz molekula H_2O i jedan iz COO grupe). Koordinacioni poliedar se može opisati kao deformisani oktaedar sa dužinama veza između 2,0550(19) i 2,1893(17) Å i uglovima u opsegu 83,06(7) – 99,52(7)°.

Tri koordinirana molekula H_2O su dupli donori vodonične veze, a jedan od njih je i akceptor. Voda kristalne rešetke je akceptor (preko O atoma iz jedne koordinirane H_2O) i donor (preko jednog O atoma iz nekoordinirane COO grupe). Postoje četiri intramolekulske vodonične veze, 2,588(3) i 2,714(2) Å, između koordiniranih H_2O i O atoma iz COO grupe.

Kristalografski podaci: $[Co_2(pyr)(phen)_2(H_2O)_6] \cdot 2H_2O$, $C_{34}H_{34}Co_2N_4O_{16}$, $M_r = 872,51$, triklinični sistem, prostorna grupa $P\bar{1}$, $a = 7,410(5)$, $b = 11,067(5)$, $c = 12,219(5)$ Å, $\alpha = 66,638(5)$, $\beta = 76,402(5)$, $\gamma = 70,855(5)$ °, $V = 862,61(8)$ Å³, $Z = 1$, $F(000) = 448$, $\rho_x = 1,680$ g cm⁻³, $\mu(Mo K\alpha) = 1,047$ mm⁻¹, 4251 izmerenih refleksija, $R_I = 3,29$ % za 3224 refleksija sa $I > 2\sigma(I)$, $R_{w2} = 7,17$ % (utačnjavanje pomoću F^2) za sve refleksije i 285 utaćnjavanih parametara, sa $(\Delta/\sigma)_{max} = 0,003$, $\Delta\rho_{max} = 0,323$ i $\Delta\rho_{min} = -0,286$ e Å⁻³.

- [1] D. Poleti, D. R. Stojaković, B. V. Prelesnik, R. M. Herak, *Acta Cryst.*, **C44** (1988), 242-245.
- [2] R.-K. Chiang, N.-T. Chuang, Ch.-Sh. Wur, M.-F. Chong, Ch.-R. Lin, *J. Solid State Chem.*, **166** (2002), 158-163.
- [3] Q. Shi, R. Cao, D.-F. Sun, M.-Ch. Hong, Y.-C. Liang, *Polyhedron*, **20** (2001), 3287-3293.

CRYSTAL STRUCTURE OF A NEW NICKEL(II) COMPLEX CONTAINING PYROMELLITATE ION AND 2,2'-DIPYRIDYLAMINE

J. Rogan^a, D. Poleti^a, Lj. Karanović^b

^a Faculty of Technology and Metallurgy, Kardex 4, Belgrade, Serbia and Montenegro,

^b Faculty of Mining and Geology, Dušina 7, Belgrade, Serbia and Montenegro

e-mail: rogan@elab.tmf.bg.ac.yu

The title binuclear complex, $[\text{Ni}_2(\text{pyr})(\text{dipy})_2(\text{H}_2\text{O})_6] \cdot 4\text{H}_2\text{O}$, is a new example of ternary transition metal complexes containing tetraanion of pyromellitic (1,2,4,5-benzenetetracarboxylic) acid, pyr, and 2,2'-dipyridylamine, dipy, as aromatic diamine ligand. The interest in such complexes is related to the numerous possible coordination modes of pyr allowing interesting double chain [1] or 3-D network [2] structures.

The structure analysis showed that Ni atoms are in a deformed octahedral environment. The dipy is a usual chelating ligand, while the pyr anion acts as a bridge and it is coordinated in a bis-monodentate manner with two uncoordinated *ortho*-COO groups. Therefore, the pyr ligand behaves in fact as a terephthalato ligand. The three remaining apices in Ni(II) octahedron are occupied by three H_2O molecules. The bond distances in the octahedron are between 2.038(3) and 2.120(2) Å and angles in the range 83.58(9) – 95.56(10)°.

The binuclear coordination unit of the title compound is very similar to the recently obtained Co(II) complex $[\text{Co}_2(\text{pyr})(\text{phen})_2(\text{H}_2\text{O})_6] \cdot 2\text{H}_2\text{O}$, phen = 1,10-phenanthroline.

Crystal packing is governed by a 3-D network of hydrogen bonds, which are in the range of 2.648(3) – 2.929(4) Å, and with O–H...O angles between 138.7(2) and 172.2(2)°. Four intramolecular H bonds between coordinated H_2O molecules and free O atoms from COO groups also exist.

Crystal data for $[\text{Ni}_2(\text{pyr})(\text{dipy})_2(\text{H}_2\text{O})_6] \cdot 4\text{H}_2\text{O}$: $C_{30}\text{H}_{40}\text{Ni}_2\text{N}_6\text{O}_{18}$, $M_r = 890.10$, triclinic, space group $P\bar{1}$, $a = 9.3420(5)$, $b = 10.8890(5)$, $c = 12.0960(5)$ Å, $\alpha = 67.382(5)$, $\beta = 76.511(5)$, $\gamma = 84.395(5)$ °, $V = 1104.5(9)$ Å³, $Z = 1$, $F(000) = 462$, $\rho_x = 1.338$ g cm⁻³, $\mu(\text{Mo K}\alpha) = 0.925$ mm⁻¹, $R_1 = 4.54\%$ for 2721 observed reflections [$I > 2\sigma(I)$], $R_{w2} = 6.95\%$ (refinement on F^2) for 4179 independent reflections and 253 refined parameters, with $(\Delta/\sigma)_{\text{max}} = 0.001$, $\Delta\rho_{\text{max}} = 0.640$ e Å⁻³, $\Delta\rho_{\text{min}} = -0.445$ e Å⁻³.

- [1] Q. Shi, R. Cao, D.-F. Sun, M.-Ch. Hong, Y.-C. Liang, *Polyhedron*, **20** (2001), 3287–3293.
- [2] W. Chen, H. H. Tioh, J.-Z. Zou, Z. Xu, X.-Z. You, *Acta Cryst.*, **C52** (1996), 43–45; Ch.-D. Wu, C.-Zh. Lu, Sh.-F. Lu, H.-H. Zhuang, J.-Sh. Huang, *Inorg. Chem. Comm.*, **5** (2002), 171–174.

KRISTALNA STRUKTURA NOVOG NIKAL(II) KOMPLEKSA SA PIROMELITAT JONOM I 2,2'-DIPIRIDILAMINOM

J. Rogan ^a, D. Poleti ^a, Lj. Karanović ^b

^a Tehnološko-metalurški fakultet, Karnegijeva 4, Beograd, Srbija i Crna Gora, ^b Rudarsko-geološki fakultet, Đušina 7, Beograd, Srbija i Crna Gora
e-mail: rogan@elab.tmf.bg.ac.yu

Navedeni binuklearni kompleks, $[Ni_2(pyr)(dipy)_2(H_2O)_6] \cdot 4H_2O$, je novi primer ternernog kompleksa prelaznih metala sa tetraanjonom piromelitne (1,2,4,5-benzentetrakarboksilne) kiseline, pyr, i 2,2'-dipiridilaminom, dipy, kao aromatičnim diaminskim ligandom. Interesovanje za ovakve komplekse potiče od brojnih mogućih načina koordinacije pyr koji dopuštaju zanimljive strukture dvostrukih lanaca [1] ili 3-D umrežene strukture [2].

Strukturalna analiza je pokazala da su atomi Ni u deformisano oktaedarskom okruženju. Dipy je uobičajeno helatni ligand, dok se pyr anjon ponaša kao most i koordiniran je na bis-monodentatni način sa dve nekoordinirane orto-COO grupe. Tako se pyr ligand praktično ponaša kao tereftalato-ligand. Ostala tri roglja u Ni(II) oktaedru zauzimaju 3 molekula H_2O . Dužine veza u oktaedru su između 2,038(3) i 2,120(2) Å, a uglovi su u opsegu 83,58(9) – 95,56(10)°.

Binuklearna kompleksna jedinka navedenog kompleksa je vrlo slična ranije dobijenom Co(II) kompleksu $[Co_2(pyr)(phen)_2(H_2O)_6] \cdot 2H_2O$, phen = 1,10-fenatrolin.

Pakovanje kristala u molekulu određeno je 3-D sistemom vodoničnih veza koje su u opsegu 2,648(3) – 2,929(4) Å i sa O–H...O uglovima između 138,7(2) i 172,2(2)°. Takođe postoje četiri intramolekulske vodonične veze između koordiniranih molekula H_2O i slobodnih O atoma iz COO grupa.


Kristalografski podaci za $[Ni_2(pyr)(dipy)_2(H_2O)_6] \cdot 4H_2O$: C₃₀H₄₀Ni₂N₆O₁₈, M_r = 890,10, triklinični sistem, prostorna grupa P $\overline{1}$, a = 9,3420(5), b = 10,8890(5), c = 12,0960(5) Å, α = 67,382(5), β = 76,511(5), γ = 84,395(5)°, V = 1104,5(9) Å³, Z = 1, F(000) = 462, ρ_x = 1,338 g cm⁻³, μ(Mo Kα) = 0,925 mm⁻¹, R₁ = 4,54 % za 2721 refleksija [$I > 2\sigma(I)$], R_{w2} = 6,95 % (utačnjavanje pomoću F²) za 4179 izmerenih refleksija i 253 utaćnjavanih parametara, sa (Δ/σ)_{max} = 0,001, Δρ_{max} = 0,640 i Δρ_{min} = -0,445 e Å⁻³.

- [1] Q. Shi, R. Cao, D.-F. Sun, M.-Ch. Hong, Y.-C. Liang, *Polyhedron*, **20** (2001), 3287-3293.
- [2] W. Chen, H. H. Tioh, J.-Z. Zou, Z. Xu, X.-Z. You, *Acta Cryst.*, **C52** (1996), 43-45; Ch.-D. Wu, C.-Zh. Lu, Sh.-F. Lu, H.-H. Zhuang, J.-Sh. Huang, *Inorg. Chem. Comm.*, **5** (2002), 171-174.

CRYSTAL STRUCTURE OF THE Ni(II) COMPLEX WITH S-METHYLISOTHOSEMICARBAZIDE AND TEREPHTHALATE

S. B. Novaković^a, G. A. Bogdanović^a, V. M. Leovac^c

^a"Vinča" Institute of Nuclear Sciences, Laboratory of Theoretical Physics and Condensed Matter Physics, 11001, Belgrade, P. O. Box 522, Serbia and Montenegro; ^bFaculty of Sciences, University of Novi Sad, 21000 Novi Sad, Serbia and Montenegro;
e-mail:snovak@vin.bg.ac.yu



Crystallographic data: $C_{12}H_{25}N_6Ni_1O_8S_2$, monoclinic system, space group $P2_1/c$, $a = 13.088(2)$, $b = 5.653(1)$, $c = 14.423(2)$ Å, $\beta = 109.55(1)^\circ$, $V = 1006.3(1)$ Å³, $Z = 2$, $D_c = 1.664$ Mg/m³, MoKα radiation, $\mu(\text{MoK}\alpha) = 1.23$ mm⁻¹, Enraf-Nonius CAD-4 diffractometer. The structure was solved using SHELXS86 and refined using SHELXL97 programs to the final R faktor 0.0438 for 1368 independent reflections with $I > 2\sigma(I)$ and 133 refined parameters.

The complex was obtained in the reaction of the aqueous solutions of $[\text{NiL}_2(\text{NO}_3)_2]$ (L = S-methylisothiosemicarbazide) and sodium-terephthalate. In the complex cation the nickel coordination geometry is distorted octahedral. The nickel atom and the terephthalate ring are positioned at symmetry centers. The crystal structure is stabilized by the complex hydrogen bonding network. The different coordination mode of S-methylisothiosemicarbazide relative to thiosemicarbazide, cause the distortion of the hydrogen bond motifs [1] usually present in the analogou complexes of thiosemicarbazide and N-substituted thiosemicarbazides [2].

[1] M. C. Etter, *J. Phys. Chem.*, 95, (1991), 4601-4610.

[2] M. T. Allen, A. D.Burrows, M. F. Mahon, *J. Chem. Soc. Dalton Trans.*, (1999), 215-221.

KRISTALNA STRUKTURA KOMPLEKSA Ni(II) SA S-METILIZOTIOSEMIKARBAZIDOM I TEREFTALATOM


S. B. Novaković^a, G. A. Bogdanović^a, V. M. Leovac^c

^aInstitut za nuklearne nauke „Vinča”, Laboratorija za teorijsku fiziku i fiziku kondenzovane materije, p. pr. 522, 11001 Beograd, Srbija i Crna Gora;

^bPrirodno-matematički fakultet,

Univerzitet u Novom Sadu, 21000 Novi Sad, Srbija i Crna Gora;

e-mali:snovak@vin.bg.ac.yu



Kristalografski podaci: $C_{12}H_{25}N_6Ni_1O_8S_2$, monoklinični kristalni sistem, prostorna grupa $P2_1/c$, $a = 13,088(2)$, $b = 5,653(1)$, $c = 14,423(2)$ Å, $\beta = 109,55(1)^\circ$, $V = 1006,3(1)$ Å³, $Z = 2$, $D_c = 1,664$ Mg/m³, MoKα zračenje, $\mu(\text{MoK}\alpha) = 1,23$ mm⁻¹. Enraf-Nonius CAD-4 difraktometar. Kristalna struktura je rešena korišćenjem programa SHELXS86 i utaćnjena programom SHELXL97 do konačnog R faktora 0,0438 za 1368 nezavisnih refleksija sa $I > 2\sigma(I)$ i 133 utaćnjениh parametra.

Kompleks je sintetisan u reakciji vodenih rastvra $[\text{NiL}_2(\text{NO}_3)_2]$ ($\text{L} = \text{S-metilizotiose-mikarbazid}$) i natrijum-tereftalata. U kompleksnom katjonu atom nikla se nalazi u deformisanom oktaedarkom okruženju i zauzima centar inverzije. Tereftalatni prsten zauzima drugi centar inverzije. Kristalna struktura je stabilizovana složenim sistemom vodoničnih veza. Razlika u koordinadiciji S-metilizotiosemikarbazida u odnosu na tiosemikarbazid uslovjava narušavanje karakterističnih motiva vodoničnih veza [1] prisutnih u analognim kompleksima tiosemikarbazida i N-metilovanih tiosemikarbazida [2].

[1] M. C. Etter, *J. Phys. Chem.*, 95 (1991), 4601-4610.

[2] M. T. Allen, A. D. Burrows, M. F. Mahon, *J. Chem. Soc. Dalton Trans.*, (1999), 215-221.

ON THE FACTORS INFLUENCING THE INTERMOLECULAR CONTACTS BETWEEN THE COPPER(II) AND PHENYL RING IN THE SQUARE-PLANAR Cu(II) COMPLEXES

Z. D. Tomić, S. B. Novaković, S. D. Zarić^a

"Vinča" Institute of Nuclear Sciences, Laboratory of Theoretical Physics and Condensed Matter Physics, 11001, Belgrade, P. O. Box 522, Serbia and Montenegro; ^aFaculty of Chemistry, Studentski trg 16, 11000 Belgrade, Serbia and Montenegro;
e-mail:zorant@vin.bg.ac.yu

There are few observations in the literature concerning the Cu(II)···C(phenyl) contacts at distances greater than 3.0 Å, and they have been discussed in terms of the bonding interaction between the copper atom and aromatic carbon. In a recent article [1] we have suggested that on the basis of the geometrical parameters and disposition of charges in the chelate and phenyl rings, this contact could be attributed to the interaction between the whole chelate ring and the phenyl fragment.

To obtain more insight into this type of interaction we have performed the analysis of the geometrical parameters of the intermolecular Cu(II)···C(phenyl) contacts in the square-planar Cu(II) complexes. Results of this investigation will be presented and the factors which could possibly influence this interaction will be discussed.

- [1] Z. D. Tomić, V. M. Leovac, S. V. Pokorni, D. Zobel, S. D. Zarić, *Eur. J. Inorg. Chem.* (2003), 1222-1226.

FAKTORI KOJI UTIČU NA INTERMOLEKULSKE KONTAKTE IZMEDJU BAKRA(II) I FENILNOG PRSTENA KOD KVADRATNO-PLANARNIH Cu(II) KOMPLEKSA

Z. D. Tomić, S. B. Novaković, S. D. Zarić^a

*Institut za nuklearne nauke „Vinča”, Laboratorija za teorijsku fiziku i fiziku kondenzovane materije, p. pr. 522, 11001 Beograd, Srbija i Crna Gora, ^aHemski fakultet, Studentski trg 16, 11000 Beograd, Srbija i Crna Gora
e-mail:zorant@vin.bg.ac.yu*

U literaturi se navodi nekoliko primera Cu(II)···C(fenil) kontakta na rastojanjima većim od 3.0 Å koji su razmatrani uz pretpostavku postojanja interakcije izmedju ova dva atoma. U našem prethodnom radu [1] ukazali smo da geometrijski parametri strukture bis[aceton-1-naftoilhidrazonato⁽⁻¹⁾]bakar(II) kompleksa, kao i raspodela naelektrisanja na atomima helatnog i fenilnog prstena sugerisu da u interakciji učestvuje na samo atom Cu već ceo helatni prsten.

Da bismo proverili osnovanost ovog zaključka izvršili smo analizu geometrijskih parametara koji opisuju Cu(II)···C(fenil) intermolekulske kontakte kod kvadratno planarnih Cu(II) kompleksa, uzimajući u obzir mogućnost da ovaj kontakt predstavlja helat-fenil interakciju. U radu će biti prikazani rezultati ove analize.

- [1] Z. D. Tomić, V. M. Leovac, S. V. Pokorni, D. Zobel, S. D. Zarić, *Eur. J. Inorg. Chem.* (2003), 1222-1226.

CRYSTAL STRUCTURE OF POTASSIUM BIS(MALONATO)CUPRATE(II) DIHYDRATE


S. R. Trifunović^a, G. N. Kaluderović^b, F. W. Heinemann^c and T. J. Sabo^b

^aDepartment of Chemistry, Faculty of Science, University of Kragujevac, 34000 Kragujevac, Serbia & Montenegro; ^bFaculty of Chemistry, University of Belgrade, Studentski trg 16, P.O. Box 158, 11001 Belgrade, Serbia & Montenegro; ^cInstitut für Anorganische Chemie II, Egerlandstrasse 1, D-91058 Erlangen, Germany
e-mail: srecko@knez.uis.kg.ac.yu

The light blue title compound was synthesised by the same procedure as for ethylenediammonium aquabis(malonato)oxovanadate(IV) [1].

Crystallographic data: $C_6H_8CuK_2O_{10}$, orthorhombic system, space group $Pbca$, $a = 7.082(1)$, $b = 9.402(1)$, $c = 16.776(1)$ Å, $V = 1117.0(2)$ Å³, $Z = 4$, $D_c = 2.271$ Mg/m³, $\mu(\text{MoK}\alpha) = 2.750$ mm⁻¹, $F(000) = 764$. Data collection: $2.4 < \theta < 26.0^\circ$, no. of parameters: 89, $R = 0.0676$, $wR_2 = 0.2136$ for 727 reflections with $I > 2\sigma(I)$.

The copper(II) is found in square-planar geometry surrounded by four carboxylate oxygen atoms from two malonato ligands. In axial position there are two carbonyl oxygen atoms, also from malonato ligands, at distance of 2.577 Å, forming distorted octahedron (angle O4–Cu–O4 180°). Around potassium atom there are six oxygen atoms, O1 and O3 which are coordinated to copper(II) ion, and O5 water molecule.



atoms, also from malonato ligands, at distance of 2.577 Å, forming distorted octahedron (angle O4–Cu–O4 180°). Around potassium atom there are six oxygen atoms, O1 and O3 which are coordinated to copper(II) ion, and O5 water molecule.

- [1] G. N. Kaluderović, F. W. Heinemann, V. M. Leovac, S. R. Trifunović, T. J. Sabo, *Acta Cryst. E59* (2003), m541-m542.

KRISTALNA STRUKTURA KALIJUM-BIS(MALONATO)KUPRAT(II) DIHIDRATA


S. R. Trifunović^a, G. N. Kaluđerović^b, F. W. Heinemann^c and T. J. Sabo^b

^aPrirodno-matematički fakultet, Univerzitet u Kragujevcu, 34000 Kragujevac, Srbija i Crna Gora; ^bHemski fakultet, Univerzitet u Beogradu, p. pr. 158, Studentski trg 16, 11001 Beograd, Srbija i Crna Gora; ^cInstitut für Anorganische Chemie II, Egerlandstrasse 1, D-91058 Erlangen, Nemačka

e-mail: srecko@knez.uis.kg.ac.yu

Svetlo plavi kompleks, $K_2[Cu(mal)] \cdot 2H_2O$ dobijen je po proceduri za etilendiammonium-akvabis(malonato)oksovanadat(IV) [1].

Kristalografski podaci: $C_6H_8CuK_2O_{10}$, ortorombični, prostorna grupa $Pbca$ $a = 7,082(1)$, $b = 9,402(1)$, $c = 16,776(1)$ Å, $V = 1117,0(2)$ Å 3 , $Z = 4$, $D_c = 2,271$ Mg/m 3 , $\mu(MoK\alpha) = 2,750$ mm $^{-1}$, $F(000) = 764$. Opseg: $2,4 < \theta < 26,0^\circ$, broj parametara: 89, $R = 0,0676$, $wR_2 = 0,2136$ za 727 refleksija sa $I > 2\sigma(I)$.



Bakar(II) jon se nalazi u kvadratno planarnoj geometriji, okružen sa četiri karboksilna kiseonikova atoma malonato liganada. U aksijalnom položaju na rastojanju od 2,577 Å nalaze se O4 karbonilni kiseonikovi atomi gradeći deformisani oktaedar oko bakar(II) jona (ugao O4–Cu–O4 180°). Kalijum(I) jon se takođe nalazi u deformisanom oktaedarskom okruženju kiseonikovih atoma i to O1 i O3, karbonilni kiseonikovi atomi malonato liganda, i O5 molekul vode.


- [1] G. N. Kaluđerović, F. W. Heinemann, V. M. Leovac, S. R Trifunović, T. J. Sabo, *Acta Cryst. E59* (2003), m541-m542.

**CRYSTAL STRUCTURE OF *TRANS*-
DIBROMO(ETHYLENEDIAMINE-*N,N'*-DI-3-
PROPIONATO)PLATINUM(IV) MONOHYDRATE**

G. N. Kaluđerović^a, D. Poletić^b, Lj. Karanović^c and T. J. Sabo^a

^aFaculty of Chemistry, University of Belgrade, P.O. Box 158, Studentski trg 16, 11001 Belgrade, Serbia & Montenegro; ^bFaculty of Technology and Metallurgy, Karnegijeva 4, 11000 Belgrade, Serbia & Montenegro ^cFaculty of Mining and Geology, Đušina 7, 11000 Belgrade, Serbia & Montenegro
e-mail: goran@chem.bg.ac.yu

The title compound was synthesised by procedure for *trans*-dichloro(ethylenediamine-*N,N'*-di-3-propionato)platinum(IV) monohydrate [1]. The research of antitumor activity, for both complexes is in progress.



Crystallographic data: PtC₈H₁₆Br₂N₂O₅.1H₂O, monoclinic system, space group P2₁/n, $a = 7.3575(8)$, $b = 11.6720(14)$, $c = 16.611(2)$ Å, $\beta = 92.70(3)^\circ$, $V = 1424.9(3)$ Å³, $Z = 4$, $D_c = 2.693$ Mg/m³, $\mu(\text{MoK}\alpha) = 1.547$ mm⁻¹, $F(000) = 1069$. Data collection: $4.0 < \theta < 35.7^\circ$, no. of parameters: 220, $R = 0.0318$, $wR_2 = 0.0619$ for 4249 reflections with $I > 2\sigma(I)$.


The platinum(IV) is hexacoordinated by two oxygen and two nitrogen atoms from eddp ligand, and two bromine atoms. It is expected that angles between Pt-Br bonds and coordinated atoms of eddp ligand are near 90°, but they show significant deviation. Namely, differences between N1-Pt-Br1 (87.6°) and N1-Pt-Br2 (94.2°) angles is about 7°. This can be explained by intramolecular interaction N1-H···Br1 ($\text{H} \cdots \text{Br}_1 = 2.638$ Å, angle N1-H···Br1 = 114.5°) and N2-H···Br2 ($\text{H} \cdots \text{Br}_2 = 2.667$ Å, angle N2-H···Br2 = 109.9°). Angle Br1-Pt-Br2 is 178.1°.

KRISTALNA STRUKTURA TRANS-DIBROMO(ETILENDIAMIN-N,N'-DI-3-PROPIONATO)PLATINA(IV) MONOHIDRATA

G. N. Kaluđerović^a, D. Poleti^b, Lj. Karanović^c and T. J. Sabo^a

^aHemijski fakultet, Univerzitet u Beogradu, P. Fax 158, Studentski trg 16, 11001 Beograd, Srbija i Crna Gora; ^bTehnološko-metalurški fakultet, Kariđegijeva 4, 11000 Beograd, Srbija i Crna Gora; ^cRudarsko-geološki fakultet, Dušina 7, 11000 Beograd, Srbija i Crna Gora
e-mail: goran@chem.bg.ac.yu

Kompleks *trans*-[Pt(eddp)Br₂] sintetisan je po postupku za dobijanje *trans*-(etilendiamin-*N,N'*-di-3-propionato)-dihloroplatina(IV) monohidrata [1] kao potencijalni antitumorski agens. Istraživanja aktivnosti oba kompleksa su u toku.



Kristalografski podaci: PtC₈H₁₆Br₂N₂O_{5,16}, monoklinični, prostorna grupa P2₁/n, $a = 7,3575(8)$, $b = 11,6720(14)$, $c = 16,611(2)$ Å, $\beta = 92,70(3)^\circ$, $V = 1424,9(3)$ Å³, $Z = 4$, $D_c = 2,693$ Mg/m³, $\mu(\text{MoK}\alpha) = 1,547$ mm⁻¹, $F(000) = 1069$. Opseg: $4,0 < \theta < 35,7^\circ$, broj parametara: 220, $R = 0,0318$, $wR2 = 0,0619$ za 4249 refleksija sa $I > 2\sigma(I)$.

Platina(IV) je heksakoordovana sa po dva atoma kiseonika i azota, iz *eddp* liganda, i dva atoma broma. Interesantno je da uglovi između Pt-Br veza i koordinovanih atoma *eddp* liganda pokazuju odstupanje od očekivanog ugla od 90° . Razlika između uglova N1-Pt-Br1 ($87,6^\circ$) i N1-Pt-Br2 ($94,2^\circ$) veća je od 7° . To se može objasniti intramolekulskim interakcijama: N1-H···Br1 ($H\cdots Br1 = 2,638$ Å, $N1-H\cdots Br1 = 114,5^\circ$) i N2-H···Br2 ($H\cdots Br2 = 2,667$ Å, $N2-H\cdots Br2 = 109,9^\circ$). Vrednost ugla Br1-Pt-Br2 je $178,1^\circ$.

[1] G. N. Kaluđerović, G. A. Bogdanović, T. J. Sabo, *J. Coord. Chem.* 55 (2002) 817-822.

SYNTHESIS AND CRUSTAL STRUCTURE OF [Pt(terpy)(cyst-S)](ClO₄)₂·0.5H₂O

T. Soldatović^a, F. W. Heinemann^b, Ž. D. Bugarčić^a

^aDepartment of Chemistry, Faculty of Science, University of Kragujevac, R. Domanovića 12, P. O. Box 60, Sr-34000 Kragujevac, Serbia and Montenegro; ^bInstitute for Inorganic Chemistry, University of Erlangen-Nürnberg, Egerlandstrasse 1, 91058 Erlangen, Germany.

e-mail: bugi@knez.uis.kg.ac.yu

The structure consists of discrete [Pt(terpy)(cyst-S)]²⁺ cation and two perchlorate ions. Terpy is coordinated to platinum as a tridentate ligand and the fourth position is occupied by L-cysteine, which is coordinated via the sulfur atom. The coordination geometry around the platinum centre is distorted square-planar with N–Pt–N angles of 80.7(3), 80.5(3) and 161.1(3) °, and N–Pt–S angles of 99.8(2), 99.0(2) and 179.6(3) °. The Pt–N(2) distance to the central nitrogen atom of the terpy ligand, 1.982(7), is shorter than those to the other two nitrogen atoms of terpy, *viz.* 2.034(7) and 2.043(7) Å. The Pt–S distance is 2.304(3) Å. All these Pt–N and Pt–S bond distances, as well as bond distances and angles of terpyridine, compare well with those reported for the similar [Pt(terpy)(thiourea)]²⁺ complex [1].

[1] Ž. D. Bugarčić, G. Liehr and R. van Eldik, *J. Chem. Soc., Dalton Trans.*, (2002), 2825 - 2830.

SINTEZA I KRISTALNA STRUKTURA [Pt(terpy)(cyst-S)](ClO₄)₂·0.5H₂O KOMPLEKSA

T. Soldatović^a, F. W. Heinemann^b, Ž. D. Bugarčić^a

^aUniverzitet u Kragujevcu, Prirodno-matematički fakultet, Institut za hemiju, R. Domanovića 12, p. pr. 60, 34000 Kragujevac, Srbija i Crna gora; ^bUniverzitet Erlangen-Nürnberg, Institut za neorgansku hemiju, Egerlandstrasse 1, 91058 Erlangen, Nemačka.
e-mail: bugi@knez.uis.kg.ac.yu

Određena je kristalna struktura kompleksa [Pt(terpy)(cyst-S)]²⁺, gde je terpy 2,2':6',2''-terpiridin, a cyst-S je L-cistein. Struktura kompleksa je kvadratno-planarna. Terpi ligand je tridentatno koordinovana, a L-cistein je koordinovan preko sumpora. Dužina Pt–N(2) veze je 1,982(7) Å (to je centralni N atom iz terpi liganda) i ta veza je kraća od druge dve Pt–N veze, 2,034(7) i 2,043(7) Å. Veza Pt–S je 2,304(3) Å, što odgovara dužini drugih Pt–S veza u sličnim kompleksima [1]. Uglovi N–Pt–N su 80,7(3), 80,5(3) i 161,1(3), a N–Pt–S su 99,8(2), 99,0(2) i 179,6(3) °. Dužine veza i uglovi u koordinovanom terpi ligandu su slične odgovarajućim vezama u kompleksu [Pt(terpy)(thiourea)]²⁺[1].

[1] Ž. D. Bugarčić, G. Liehr and R. van Eldik, *J. Chem. Soc., Dalton Trans.*, (2002), 2825 – 2830.

SYNTHESIS AND CRUSTAL STRUCTURE OF [Pd(terpy)(py)](ClO₄)₂

B. Petrović^a, E. Zangrandeo^b and Ž. D. Bugarčić,^a

^a*University of Kragujevac, Faculty of Science, Department of Chemistry, R. Domanovića 12, P. O. Box 60, Sr-34000 Kragujevac, Serbia and Montenegro;* ^b*University of Trieste, Department of Chemistry, I-34127 Trieste, Italy.*

e-mail: bugi@knez.uis.kg.ac.yu

The structure consists of discrete [Pd(terpy)(py)]²⁺ cation and two perchlorate ions. The metal coordination sphere is square planar with expected deviations from right angles about Pd. The Pd–N bond length to the central atom of terpy ligand is well below 2.0 Å and it is significantly shorter than any of the other M–N distances (Table 1). The terpy ligand is almost coplanar with maximum displacement from the mean plane of -0.124(4) and +0.146(6) Å exhibited by N(1) and C(3), respectively. The pyridine forms a dihedral angle of 61.9(2) ° with the coordination N4 donors. These geometrical features are comparable to those observed in the related species containing methylcytosine [1].

The crystal packing evidences perchlorate anions located upper and below the metal (the shortest metal–oxygens contacts are about 3.48 Å), and the pyridine rings of centrosymmetric pairs of complexes that are marginally stacked, the shorter C–C distance being 3.5 Å.

Table 1. Selected bond distances (Å) and angles (°).

Pd–N(1)	2.039(4)	Pd–N(3)	2.038(4)
Pd–N(2)	1.932(4)	Pd–N(4)	2.038(4)
N(1)–Pd–N(2)	81.18(17)	N(2)–Pd–N(3)	80.72(17)
N(1)–Pd–N(3)	161.70(17)	N(2)–Pd–N(4)	177.48(16)
N(1)–Pd–N(4)	97.97(17)	N(3)–Pd–N(4)	100.22(16)

- [1] S. Cosar, M. B. L. Janik, M. Flock, E. Freisinger, E. Farkas, B. Lippert, *J. Chem. Soc., Dalton Trans.*, (1999), 2329-2333.

SINTEZA I KRISTALNA STRUKTURA [Pd(terpy)(py)](ClO₄)₂ KOMPLEKSA

B. Petrović^a, E. Zangrando^b i Ž. D. Bugarčić,^a

^a Univerzitet u Kragujevcu, Prirodno-matematički fakultet, Institut za hemiju R. Domanovića 12, p. pr. 60, Sr-34000 Kragujevac, Srbija i Crna Gora; ^b Univerzitet u Trstu, Institut za hemiju, I-34127 Trst, Italija.

e-mail: bugi@knez.uis.kg.ac.yu

Određena je struktura [Pd(terpy)(py)]²⁺ kompleksa, gde je terpy 2,2': 6',2''-terpiridin, a py je piridin. Kompleks je kvadratno-planaran. Terpi je tridentatno koordinovan, a četvrto mesto zauzima N iz piridina. Dužina Pd–N(2) veze (centralni atom N u terpi ligandu) je kraća od druge dve Pd–N veze, Tabela 1.

Table 1. Dužine veza (Å) i uglovi (°) u [Pd(terpy)(py)]²⁺ kompleksu.

Pd–N(1)	2,039(4)	Pd–N(3)	2,038(4)
Pd–N(2)	1,932(4)	Pd–N(4)	2,038(4)
N(1)–Pd–N(2)	81,18(17)	N(2)–Pd–N(3)	80,72(17)
N(1)–Pd–N(3)	161,70(17)	N(2)–Pd–N(4)	177,48(16)
N(1)–Pd–N(4)	97,97(17)	N(3)–Pd–N(4)	100,22(16)

Dobijeni rezultati se mogu porebiti sa ranije objavljenom strukturom paladijum kompleksa [1].

- [1] S. Cosar, M. B. L. Janik, M. Flock, E. Freisinger, E. Farkas, B. Lippert, *J. Chem. Soc., Dalton Trans.*, (1999), 2329-2333.

BaMgAs₂O₇ AND BaCoAs₂O₇ – SYNTHESIS AND CRYSTAL STRUCTURES OF TWO NEW BARIUM DIARSENATES

T. Mihajlović, H. Effenberger and E. Tillmanns

*Institut für Mineralogie und Kristallographie, Universität Wien – Geozentrum,
Althanstrasse 14, 1090 Wien, Austria.*

e-mail: tamara.mihajlovic@univie.ac.at

During work aimed at the hydrothermal preparation of synthetic members of the descloizite-adelite group, the title compounds were prepared by hydrothermal methods. Mixtures of Ba(OH)₂, MgO/Co(OH)₂, H₃AsO₄ and H₂O were transferred into teflon-lined stainless steel autoclaves and heated up to 493 K. The reactions were maintained under autogeneous pressure for three days followed by slow cooling to room temperature. The final products were filtered and washed thoroughly with distilled water. BaMgAs₂O₇ crystallised as transparent colourless prismatic crystals up to 0.15 mm in length and BaCoAs₂O₇ as transparent blue-green crystals up to 0.19 mm in length. These diarsenates represent two new members belonging to the M₁²⁺M₂²⁺As₂O₇ family of compounds.

Prior to our study, only five diarsenates with the general formula M₁²⁺M₂²⁺As₂O₇ were known [1]. Four of these (PbCuAs₂O₇, BaCuAs₂O₇, SrCuAs₂O₇ and SrCoAs₂O₇) crystallise in space group *P*2₁/*n* with *Z* = 4, and the fifth one (CaCuAs₂O₇) crystallise in *P*2₁/*c*, but has different structure. The structures of these compounds consist of corner-sharing M₂O₅ square pyramids and As₂O₇ diarsenate groups, where M₁ atoms occupy the gap between two layer-like [M₂As₂O₇] units.

The structures of the title compounds were solved by direct methods [2] from single-crystal X-ray diffraction data (298 K, CCD detector, MoK α , $2\theta_{\max} = 65^\circ$) obtained from non-merohedrally twinned crystals [twin plane (100)]. The structure models were refined using SHELXL-97 [3]. The space group symmetry of both diarsenates is *P*2₁/*n* with *Z* = 4. BaMgAs₂O₇ has *a* = 5.620(1), *b* = 8.629(2), *c* = 13.344(3) Å, β = 90.20(3) $^\circ$, *V* = 647.1(2) Å³ with *R*₁ = 2.42 % (*wR*₂ = 5.26%) for 2349 unique reflections with, and *R*₁ = 2.25% for 2269 observed reflections using 102 free parameters. BaCoAs₂O₇ has *a* = 5.651(1), *b* = 8.580(2), *c* = 13.282(3) Å, β = 90.01(3) $^\circ$, *V* = 664.0(2) Å³ with *R*₁ = 2.21% (*wR*₂ = 4.99%) for 2347 unique reflections with, and *R*₁ = 2.09 % for 2286 observed reflections using 102 free parameters. The structures are isotopic with the other M₁²⁺M₂²⁺As₂O₇ diarsenates crystallising in *P*2₁/*n*. The M₂ position is five coordinated, which is rare coordination for Co and very rare for Mg.

Financial support of the Austrian Science Foundation (FWF) (Grant P15875-N03 to H. Effenberger) is gratefully acknowledged.

- [1] T. C. Chen and S. L. Wang, *J. Solid State Chem.*, 121 (1996), pp 350-355.
- [2] G. M. Sheldrick, SHELXS-97, a program for the solution of crystal structures. University of Göttingen, Germany (1997).
- [3] G. M. Sheldrick, SHELXL-97, a program for crystal structure refinement. University of Göttingen, Germany (1997).

BaMgAs₂O₇ AND BaCoAs₂O₇ – SINTEZA I KRISTALNA STRUKTURA DVA NOVA DIARSENATA BARIJUMA

T. Mihajlović, H. Effenberger i E. Tillmanns

*Institut für Mineralogie und Kristallographie, Universität Wien – Geozentrum,
Althanstrasse 14, 1090 Wien, Austria.*

e-mail: tamara.mihajlovic@univie.ac.at

U toku hidrotermalne sinteze novih sintetičkih članova grupe desklozit-adelit, kristalisali su novi diarsenati barijuma. Za sintezu je korišćena mešavina Ba(OH)₂, MgO/Co(OH)₂, H₃AsO₄ i H₂O tretirana u teflonima zagrevanim u anklavama od nerđajućeg čelika na 493 K. Reakcija se odigravala pod autogenim pritiskom 3 dana i praćena je sporim hlađenjem do sobne temperature. Finalni produkti su filtrirani i detaljno oprani destilovanom vodom. BaMgAs₂O₇ je iskristalisaao kao providan, bezbojan, prizmatični kristal dužine do 0,15 mm, a BaCoAs₂O₇ kao providan, plavo-zeleni kristal dužine do 0,19 mm. Ovi diarsenati predstavljaju dva nova člana familije supstanci sa opštom formulom M₁²⁺M₂²⁺As₂O₇.

Do sada je bilo poznato samo pet M₁²⁺M₂²⁺As₂O₇ diarsenata [1]. Četiri od njih (PbCuAs₂O₇, BaCuAs₂O₇, SrCuAs₂O₇ i SrCoAs₂O₇) kristališu u prostornoj grupi P₂₁/n sa Z = 4, a peti (CaCuAs₂O₇) kristališe u P₂₁/c i pokazuje drugačiju strukturu. Struktura ovih stupstanci sastoji se od M₂O₅-piramide povezanih rogljevima sa As₂O₇-diarsenat grupama i M₁ atoma koji se nalaze u kanalima između dva sloja [M₂As₂O₇]-jedinica.

Struktura diarsenata barijuma rešena je pomoću direktnih metoda [2] korišćenjem podataka dobijenih rendgenskom difrakcijom na monokristalu (298 K, CCD detektor, MoKα, 2θ_{max} = 65°) i to nemeroherdalno bližnjeni kristala [ravan bliznjenja je (100)] u oba slučaja]. Strukturni model utaćnjavan je pomču SHELLXL [3]. Oba diarsenata su utaćnjena u prostornoj grupi P₂₁/n sa Z = 4. Parametri jedinične celije za BaMgAs₂O₇ su: a = 5,620(1), b = 8,629(2), c = 13,344(3) Å, β = 90,20(3)°, V = 647,1(2) Å³, a R₁ = 2,42% (wR₂ = 5,26%) za 2349 nezavisnih refleksija i R₁ = 2,25% za 2269 primećenih refleksija i 102 utaćnjavana parametra. Parametri jedinične celije za BaCoAs₂O₇ iznose a = 5,651(1), b = 8,580(2), c = 13,282(3) Å, β = 90,01(3)°, V = 664,0(2) Å³ sa R₁ = 2,21% (wR₂ = 4,99%) za 2347 nezavisnih refleksija i R₁ = 2,09% za 2286 primećenih refleksija za 102 slobodno utaćnjavana parametra. Oba barijum diarsenata su izostrukturna sa ostalim M₁²⁺M₂²⁺As₂O₇ diarsenatima koji kristališu u prostornoj grupi P₂₁/n. Zanimljivo je napomenuti da M₂ katjon kod M₁²⁺M₂²⁺As₂O₇ diarsenata pokazuje koordinaciju pet, što je retko okruženje za Co, a izuzetno retko okruženje za magnezijum.

Zahvaljujemo se Austrijskom fondu za nauku (FWF) (projekat P15875-N03 na ime H. Effenberger).

[1] T. C. Chen and S. L. Wang, *J. Solid State Chem.*, 121 (1996), pp 350-355.

[2] G. M. Sheldrick, SHELLXS-97, a program for the solution of crystal structures.

University of Göttingen, Germany (1997).

[3] G. M. Sheldrick, SHELLXL-97, a program for crystal structure refinement. University of Göttingen, Germany (1997).

THE STATE OF ART AND EXPECTED DEVELOPMENTS ON THE CALIBRATION OF DIFFRACTOMETER

Giovanni Berti

University of Pisa – Via S.Maria 53, 56126 Pisa - Italy
 e-mail: g.bert@ing.unipi.it

The calibration of x-ray diffractometers implies to evaluate the residual systematic effects that are still present after the instrument alignment. When the alignment is performed as better as possible residual and not removable contributions to the diffraction pattern are present and their quantification possible. This evaluation is carried out by calculating the effective values of the parameters involved in the whole diffraction measurement process. These effective values are related to the actual experimental conditions, rather than the nominal values of the individual devices. The nominal values are provided by the manufacturers after the test on its site. The effective values of parameters involved in the diffraction experiment, and the effective functioning of the individual devices at the user site; these effective values enable the visualisation of “characteristic curves” of the diffraction measurements. When the uncertainty is associated to these curves, a reasonable confidence of the measurement reproducibility and repeatability may be reached.

The calibration process requires that the basic principle of the diffraction physics are considered. We report here the results obtained by using the Diffraction Instrumental Monitoring Methods and the protocol used by DISVAR96 package. In principle it can be extended to other packages and protocols useful for the diffraction pattern interpretation. Other developments are expected by extending the theory developed by Wilson [1]. This theory is in fact sometimes applied to divergent beams, where the gross effects (zero shift, divergence – axial and equatorial, specimen surface displacements) usually cover any other systematic effect. There are anyway some other secondary effects that become relevant when the gross effects are negligible as in the case of parallel beam geometry. These effects (lorentz, wavelength dispersion and even refraction) can contribute to the line position shift and the line broadening together other effects (2:1 missetting, inclination of the goniometer axis etc.).

One of the crucial points is the verification of the internal consistency of the parameters related to profile analysis of the diffraction lines. This internal consistency verification concerns the dependence on 2θ of the line position, the line broadening (FWHM), the mixing parameter (when the pseudo-Voigt is used) the normalised intensity. The fine analysis of this consistency requires that the verification will be extended to the asymmetric contribution to the diffraction line. When this asymmetric contribution can be analysed as a lorentzian addition to the profile analysis, the result is shown by the following Figures (Figs. 1 and 2).

The second crucial point is the horizontal and vertical balance of the table reporting the effective values of the individual devices contributing to the systematic

effects. The vertical balance depends on individual diffractometers, the horizontal balance involves all the diffractometers participating to the calibration net.




Figure 1. Characteristic curves with correspondent errors for the diffractometers A (up) and B (down); solid line (first cycle), dash line (second cycle) and dot-dash line (third cycle).




Figure 2. Evaluation of internal consistency of data for calibration purposes. When the Lorentzian function is used to represent the asymmetry the parameter I_3 represents is intensity of the aberration and I is the intensity of the $CuK\alpha_1$ radiation. Si_3 is the FWHM of asymmetry.

- [1] A.J.C. Wilson (1963) "The Mathematical Theory of Powder Diffraction" Philips Technical Library.
- [2] G. Berti, U. Bartoli, M. D'Acunto, F. De Marco, *Material Science Forum*, in press.
- [3] G. Berti, *Powder Diffraction* 16 (2001), p. 1-5
- [4] G. Berti, *Powder Diffraction*, 16 (2001), p. 6-15.

STRUCTURAL PROPERTIES OF $\text{La}_{1-x}\text{Na}_x\text{MnO}_3$ SOLID SOLUTIONS

M. Mitrić ^a, J. Blanuša ^a, Z. Dohčević – Mitrović ^b, N. Paunović ^b, V. Spasojević ^a

^a Institute for Nuclear Sciences "Vinča", P. O. Box 522, 11001 Belgrade; Serbia and Montenegro, ^b Institute of Physics, P. O. Box 57, 11001 Belgrade, Serbia and Montenegro
e-mail: jblanusa@ptt.yu

$\text{La}_{1-x}\text{Na}_x\text{MnO}_3$ ($x = 0; 0.03; 0.06; 0.09; 0.12; 0.15$) solid solutions were synthesized by ceramic technology.

The X-ray powder diffraction data were collected on a Philips PW1050 powder diffractometer with Ni-filtered CuK_α radiation. The data were collected in the 2θ region $10\text{--}110^\circ$, with a step size of 0.02° and an exposure time of 10 s per step. The crystal structures were refined with the use of the Fullprof software package.

It was found that all samples crystallize in trigonal structure, in the space group R-3c. Also, it was found that after Glazer notation the configuration of oxygen octahedra belongs to $\text{a}^- \text{a}^- \text{a}^-$ type [1]. It is shown that the tilt angle between oxygen octahedra lies within $15\text{--}18^\circ$ and it increases with the sodium content.

- [1] A. M. Glazer, *Acta Cryst.*, B28 (1972), p. 3384-3392.

STRUKTURNЕ OSOBINE ČVRSTIH RASTVORA $\text{La}_{1-x}\text{Na}_x\text{MnO}_3$

M. Mitrić ^a, J. Blanuša ^a, Z. Dohčević – Mitrović ^b, N. Paunović ^b, V. Spasojević ^a

^a Institut za nuklearne nauke „Vinča”, p. pr. 522, 11000 Beograd, Srbija i Crna Gora;

^b Institut za fiziku u Zemunu, p. pr. 57, 11001 Beograd, Srbija i Crna Gora

e-mail: jblanusa@ptt.yu

Čvrsti rastvori $\text{La}_{1-x}\text{Na}_x\text{MnO}_3$ ($x = 0, 0,03; 0,06; 0,09; 0,12; 0,15$) sintetisani su klasičnom keramičkom tehnologijom.

Difraktogrami uzoraka snimljeni su na difraktometru za prah Philips PW1050 sa bakarnom anodom i Ni-filterom. Uzorci su snimani u intervalu $10 - 110^\circ 2\theta$ sa korakom od $0,02^\circ$ i vremenom ekspozicije od 10 s. Za utačnjavanje kristalne strukture korišćen je program Fullprof.

Nadjeno je da svi uzorci kristališu u trigonalnoj prostornoj grupi R-3c, kao i da je konfiguracija zaokretanja kiseoničnih oktaedara u ovom slučaju tipa a⁻a⁻a⁻ po Glazer-ovoј notaciji [1]. Pokazano je da se ugao zakretanja kiseoničnih oktaedara kreće od $15-18^\circ$ i da se povećava sa koncentracijom natrijuma.

[1] A. M. Glazer, *Acta Cryst.*, B28 (1972), str. 3384-3392.

DETERMINATION OF MINERAL COMPOSITION OF URINARY CALCULI BY X-RAY POWDER DIFFRACTION ANALYSIS

A. Rosić^a, S. Radulović^b

^aFaculty of Mining and Geology, Đušina 7, Belgrade, Serbia and Montenegro; ^bKBC Zvezdara, Dimitrija Tucovića 161, Belgrade, Serbia and Montenegro
e-mail: srsasa@afrodita.rcub.bg.ac.yu

Urinary calculi are crystals formed mainly of one or more of the three components, oxalate (Whewellite - $\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$, Weddellite - $\text{CaC}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$), phosphate (Hydroxylapatite - $\text{Ca}_5(\text{PO}_4)_3(\text{OH})$, Whitlockite - $\text{Ca}_5(\text{PO}_4)_3$) and urate (Uricite - $\text{C}_4(\text{NH})_2\text{O}_2\text{C}(\text{NH})_2\text{O}$). These stones may contain various combination of cations, but the most common type of stone contains calcium in combination with either oxalate or phosphate. Cystine stones (*l*-Cystine - $\text{C}_6\text{H}_{12}\text{N}_2\text{O}_4\text{S}_2$) are less common. Urinary stones have become increasingly common in most parts of the world. [1]

The analysis of the urinary calculi only by chemical methods are rather unsatisfactory [2]. X-ray diffraction identifies the constituent of a calculus by their unique diffraction patterns, which allowes definite identification of an unknown crystalline substance. The mayor advantage of X-ray diffraction is it's almost absolute identification of crystalline materials and mixtures of crystalline materials.

More then fifty samples of urocalculi from one of the Belgrade hospitals are investigated in this work. The results of X-ray diffraction analysis indicated that most of these samples belong to the oxalate and mixed type of oxalate-phosphate stones and represent up to 70 % of the investigated calculi. It has been shown that the mineral composition of the urocalculi relates to patient age.

- [1] S. Nasir *et al.*, *Oxford Research Forum*, On-line journal., Art. 5.
- [2] M. Daudon, R. J. Reveillaud, *Presse Med.*, (1987), Apr. 11, 16 (13), 627-631.

ОДРЕЂИВАЊЕ МИНЕРАЛНОГ САСТАВА УРИНАРНОГ КАМЕЊА МЕТОДОМ РЕНДГЕНСКЕ ДИФРАКЦИЈЕ ПРАХА

A. Росић^a, С. Радуловић^б

^a Рударско-геолошки факултет, Ђушина 7, Београд, Србија и Црна Гора; ^б КБЦ Звездара, Димитрија Туцовића 161, Београд, Србија и Црна Гора,
e-mail: srsasa@afrodita.rcub.bg.ac.yu

Уринарно камење чине кристали настали углавном од једне или више компоненти: оксалата (вејелит - $\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$, веделит - $\text{CaC}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$), фосфата (хидроксиапатит - $\text{Ca}_5(\text{PO}_4)_3(\text{OH})$, вајлокит - $\text{Ca}_5(\text{PO}_4)_3$) и урата (урицит - $\text{C}_4(\text{NH})_2\text{O}_2\text{C}(\text{NH})_2\text{O}$). Могу да садрже разне катјоне, али најчешће садрже калцијум у комбинацији са оксалат- или фосфат-јонима. Камење изграђено од цистина (*l*-цистин - $\text{C}_6\text{H}_{12}\text{N}_2\text{O}_4\text{S}_2$) се ређе јавља. Појава уринарног камења је све заступљенија у многим деловима света [1].

Испитивање уринарног камења само хемијским методама често није задовољавајуће [2]. Дифракција рендгенских зрака идентификује минерални састав камена и обезбеђује потпуно дефинисање непознате кристалне супстанце. Основна предност примене методе рендгенске дифракције јесте у могућности идентификације како једне тако и мешавине више кристалних супстанци.

У овом раду испитано је више од 50 узорака уринарног камења пацијената из једне београдске болнице. Резултати рендгенских испитивања указују да већина узорака припада оксалатима или мешаном типу оксалат - фосфат, што представља више од 70 % испитиваних узорака. Показана је и зависност минералног састава уринарног камења у односу на старост пацијената.

[1] S. Nasir *et al.*, *Oxford Research Forum*, On-line journal., Art. 5.

[2] M. Daudon, R. J. Reveillaud, *Presse Med.*, (1987), Apr. 11, 16 (13), 627-631.

STRUCTURE AND MICROSTRUCTURE OF CATION DEFICIT Zn-Mn FERRITES

A. Kremenović^a, B. Antić^b, A. S. Nikolić^c and M. Stoiljković^d

^a Faculty of Mining and Geology, Laboratory for Crystallography, University of Belgrade, P.O. Box 162, 11001 Belgrade, Serbia and Montenegro; ^b Institute of Nuclear Sciences "Vinča", Laboratory of Solid State Physics, P.O. Box 522, 11001 Belgrade, Serbia and Montenegro; ^c Faculty of Chemistry, Department for Inorganic Chemistry, University of Belgrade, P.O. Box 158, 11001 Belgrade, Serbia and Montenegro; ^d Institute of Nuclear Sciences "Vinča", Laboratory of Physical Chemistry, P.O. Box 522, 11001 Belgrade, Serbia and Montenegro
e-mail: akremen@eunet.yu

The ultrafine ZnFe₂O₄, MnFe₂O₄ and cation deficit Zn-Mn ferrites were obtained by thermal decomposition of appropriate mixed complex compounds with acetylacetone(2,4-pentadione) ligands ([M(AA)_x]) at 500 °C. In ZnFe₂O₄ cation distribution is partial inverse, with about 20% of Zn²⁺ ions at octahedral 16d sites, while MnFe₂O₄ is a normal spinel. Cation distribution in nonstoichiometric (Zn,Mn,Fe)_{3-x}O₄ ($\delta \approx 0.28-0.34$) is found to be (Zn_xMn_y)₈[Fe₂]_v_{16d}O₄, with a random distribution of vacancies. Nonstoichiometry in Zn-Mn ferrites is accompanied with the cation valence change, *i.e.* partial oxidation of Mn²⁺ to Mn⁴⁺. Microstructure size-strain analysis [1,2] shows that the crystallite size decreases while crystallite strain increases with the Mn-concentration increase. Three-dimensional representation of crystallite size and strain by cubic harmonic function indicates that for ZnFe₂O₄ average crystallite is in the "rhombohedron like" form while strains are higher in the crystal axes directions than between them. As the Mn-concentration increases this trend is less pronounced and three-dimensional body looks more like a spheroid.

- [1] D. Balzar, *J. Res. Natl. Inst. Stand. Technol.*, **98** (1993), 321.
- [2] J. Rodriguez-Carvajal, FullProf computer program, 1998,
<ftp://charybde.saclay.cea.fr/pub/divers/fullprof.98/windows/winfp98.zip>

STRUKTURA I MIKROSTRUKTURA KATJONSKI DEFICITARNIH Zn-Mn FERITA

A. Kremenović^a, B. Antić^b, A. S. Nikolić^c i M. Stojljković^d

^a Rudarsko geološki fakultet, Laboratorija za kristalografsku Univerzitet u Beogradu, p. pr. 162, 11001 Beograd, Srbija i Crna Gora; ^b Institut za nuklearne nauke „Vinča”,

Laboratorija za fiziku čvrstog stanja, p. pr. 522, 11001 Beograd, Srbija i Crna Gora;

^c Hemijski fakultet, Katedra za neorgansku hemiju, Univerzitet u Beogradu, p. pr. 158, 11001 Beograd, Srbija i Crna Gora; ^d Institut za nuklearne nauke „Vinča”, Laboratorija za fizičku hemiju, p. pr. 522, 11001 Beograd, Srbija i Crna Gora

e-mail: akremen@eunet.yu

Ultrafini prahovi $ZnFe_2O_4$, $MnFe_2O_4$ i katjonski deficitarnih Zn-Mn ferita dobijeni su termalnim razlaganjem odgovarajućih mešovitih kompleksnih jedinjenja sa acetilaceton(2,4-pentadijon) ligandima ($[M(AA)_x]$) na $500\ ^\circ C$. Katjonska raspodela u $ZnFe_2O_4$ je delimično inverzna, sa oko 20% Zn^{2+} jona u oktaedarskim $16d$ mestima, dok je $MnFe_2O_4$ spinel sa normalnom raspodelom katjona. Katjonska raspodela u nestehiometrijskim $(Zn,Mn,Fe)_{3-\delta}O_4$ ($\delta \approx 0.28-0.34$) je $(Zn_xMn_y)_{8a}[Fe_{z-l-v}]_{16d}O_4$, sa slučajnom raspodelom vakancija (\square). Nestehiometrija u Zn-Mn feritima povezana je sa promenom valentnog stanja, tj. parcijalnom oksidacijom Mn^{2+} u Mn^{4+} . Analiza mikrostrukture [1,2] pokazuje da se dimenzija kristalita smanjuje a naprezanje u kristalitu povećava sa povećanjem koncentracije mangana. Trodimenzionalni prikaz kristalita i naprezanja u kristalitu za $ZnFe_2O_4$ pomoću kubne harmonijske funkcije ukazuje da je kristalit romboedarskog oblika i da je naprezanje izraženije u pravcu kristalografskih osa nego između njih. Sa povećanjem koncentracije mangana ovaj trend je sve manje izražen a trodimenzionalno telo sve više ima formu sferoida.

[1] D. Balzar, *J. Res. Natl. Inst. Stand. Technol.*, **98** (1993), 321.

[2] J. Rodriguez-Carvajal, FullProf computer program, 1998,

<ftp://charybde.saclay.cea.fr/pub/divers/fullprof.98/windows/winfp98.zip>

MICROSTRUCTURE PARAMETERS OF Na – NEPHELINES WITH DIFFERENT Si/Al RATIO

P. Vulić^a, R. Dimitrijević^a, V. Dondur^b, S. Marković^b

^a Department of Crystallography, Faculty of Mining and Geology, Đušina 7, 11000 Belgrade, Serbia and Montenegro; ^b Faculty of Physical Chemistry, Studentski trg 12-16, P.O. Box 137, 11000 Belgrade, Serbia and Montenegro
e-mail: pvulic@mkgf.rgf.bg.ac.yu

Nepheline $(K,Na)SiAlO_4$ is mineral from feldspatoide group. It is stuffed derivative of tridymite structure, where interstitial positions are occupied with sodium and potassium. In nature, up to one quarter of those positions are occupied by potassium. Applying zeolite thermally induced transformation three samples of sodium nephelines were synthesized. Depending on chemical composition of starting zeolites: Na-LTA (zeolite 4A, Si/Al = 1.00), Na-FAU (zeolite 13X, Si/Al = 1.24) and Na-GIS (zeolite P, Si/Al = 1.50), one stoichiometric and two nonstoichiometric nephelines were obtained [1]. For these three samples microstructure parameters are calculated by the Warren – Averbach method, in order to obtain dependence between microstructure and Si/Al ratio. Results show increase of crystallite size and decrease of microstrain with disordering of framework cations.

- [1] P. Vulić, R. Dimitrijević, V. Dondur, S. Macura, VIII Conference of the Serbian Crystallographic Society, Book of Abstracts, Kragujevac 2000, p. 77.

МИКРОСТРУКТУРНИ ПАРАМЕТРИ Na – НЕФЕЛИНА СА РАЗЛИЧИТИМ ОДНОСОМ Si/Al

П. Вулић^a, Р. Димитријевић^a, В. Дондур^b, С. Марковић^b

^a Катедра за кристалографију, Рударско – геолошки факултет, Ђушина 7, 11000 Београд, Србија и Црна Гора; ^b Факултет за физичку хемију, Студентски трг 12-16, ПП 137, 11000 Београд, Србија и Црна Гора;

е-пошта: pvulic@mkgp.rgf.bg.ac.yu

Нефелин $(\text{K}, \text{Na})\text{SiAlO}_4$ је минерал из групе фелдспатоида. Његова структура може се извести из структуре тридимита, попуњавањем интерстицијских положаја натријумом и калијумом. У природном минералу калијум заузима до једне четвртине интерстицијских положаја. Применом термално индуковане трансформације зеолита раније су синтетизована три узорка натријумских нефелина. У складу са хемијским саставом полазних зеолита: Na-LTA (зеолит 4A, Si/Al = 1,00), Na-FAU (зеолит 13X, Si/Al = 1,24) и Na-GIS (зеолит P, Si/Al = 1,50), добијени су један стехиометријски и два не-стехиометријска нефелина [1]. За сва три узорка одређени су микроструктурни параметри Warren – Averbach-овом методом, да би се успоставила зависност између микроструктуре и односа Si/Al. Резултати указују на повећање величине кристалита и смањење микронапрезања са смањењем уређености мрежних катјона.

- [1] P. Vulić, R. Dimitrijević, V. Dondur, S. Macura, VIII Conference of the Serbian Crystallographic Society, Book of Abstracts, Kragujevac 2000, p. 77.

2Bi₂O₃·3TiO₂ SYSTEM: STRUCTURAL CHANGES INDUCED BY INTENSIVE MILLING

M. Zdujić ^a, Č. Jovalekić ^b, Lj. Karanović ^c, D. Poleti ^d

^a*Institute of Technical Sciences of SASA, Belgrade, Knez Mihailova 35;* ^b*Center for Multidisciplinary Studies, University of Belgrade, Kneza Višeslava 1, Belgrade;* ^c*Faculty of Mining and Geology, Dušina 7, Belgrade;* ^d*Faculty of Technology and Metallurgy, Kardeljeva 4, Belgrade;*
e-mail: zdujic@itn.sanu.ac.yu

Previously, we have shown that mechanochemical treatment of either 2Bi₂O₃·3TiO₂ mixture or Bi₄Ti₃O₁₂ compound leads to a gradual formation of an amorphous phase, regardless of the initial state [1]. Here we report an attempt to derive relevant structural parameters: crystallite size, amount of amorphous phase and the transformed fraction (as a result of chemical reaction between Bi₂O₃ and TiO₂) of the powders milled for various milling times. The XRPD spectra are deconvoluted with the aid of Renex and PeakFit programs. The spectra were modelled as an amorphous halo plus a set of pseudo-Voigt peaks. The fraction of amorphous phase is estimated by the ratio of the integrated diffraction intensity of the amorphous halo to the total integrated diffraction intensity in the range 22.5–36.5 °2θ. For this procedure the calibration curve has been prepared using a series of mixtures of crystalline and "fully" amorphous Bi₄Ti₃O₁₂ powders in various proportions, *i.e.* 0, 20, 40, 50, 60, 80 and 100 wt.% of the crystalline phase.

The obtained structural parameters were used to follow kinetics of the processes taking place during mechanochemical treatment: crystallite size reduction, formation of amorphous phase and chemical reaction.

From such structural analysis it was concluded that amorphisation of the Bi₄Ti₃O₁₂ compound starts when crystallite size decreases to some critical value (below 400 Å). As the formation of amorphous phase proceeds, crystallite size further slightly decreases. In the Bi₂O₃–TiO₂ mixture, in which mechanically induced chemical reaction occurs, nanocrystalline Bi₄Ti₃O₁₂ was formed within the milling period up to 1 h. As chemical reaction continues, initially formed nanocrystalline phase transforms to the amorphous phase.

It was demonstrated that steady state (nanocrystalline/amorphous phase ratio) depends on milling conditions: higher milling intensity induces "nanocrystallisation" of amorphous phase, *i.e.* precipitation of crystallites with the average size of about 170 Å.

- [1] M. Zdujić, Č. Jovalekić, Lj. Karanović, D. Poleti, X Conference of the Serbian Crystallographic Society, Book of Abstracts, Belgrade 2002, p. 37.

2Bi₂O₃·3TiO₂ СИСТЕМ: СТРУКТУРНЕ ПРОМЕНЕ ИНДУКОВАНЕ ИНТЕНЗИВНИМ МЛЕВЕЊЕМ

М. Здујић^a, Ч. Јовалекић^b, Љ. Карановић^b, Д. Полети^c

^a Институт техничких наука САНУ, Кнез Михаилова 35, Београд; ^b Центар за мултидисциплинарне студије, Кнеза Вишеслава 1, Београд; ^c Рударско-геолошки факултет, Ђушина 7, Београд; ^c Технолошко-металурички факултет, Карнегијева 4, Београд;
е-пошта: zdujic@itn.sanu.ac.yu

Раније је показано да механохемијски третман смеше састава 2Bi₂O₃·3TiO₂ или одговарајућег једињења Bi₄Ti₃O₁₂ води до постепеног образовања аморфне фазе, без обзира на различита почетна стања [1]. У овоме раду учињен је покушај да се у праховима млевеним различито време одреде релевантни структурни параметри: величина кристалита, количина аморфне фазе и количина добијеног производа (као последице хемијске реакције између Bi₂O₃ и TiO₂). Извршена је деконволуција пикова на дифрактограмима прахова помоћу програма Renex и PeakFit. Спектри су описивани аморфним брегом и пиковима типа pseudo-Voigt. Удео аморфне фазе процењен је из односа интегралног дифрактованог интензитета аморфног брега према укупном интегралном дифрактованом интензитету у опсегу 22,5–36,5 °2θ. За ову сврху направљена је калибрациона крива од смеша кристалног и „потпуно“ аморфног праха Bi₄Ti₃O₁₂ различитог састава и то: 0, 20, 40, 50, 60, 80 и 100 мас. % кристалне фазе.

Добијени структурни параметри коришћени су за праћење кинетике процеса који се одигравају током механохемијског третмана: смањење величине кристалита, образовање аморфне фазе и хемијска реакција.

Из ове структурне анализе закључено је да аморфизација Bi₄Ti₃O₁₂ једињења отпочиње тек након смањења кристалита до неке критичне величине (испод 400 Å). У наставку образовања аморфне фазе величина кристалита споро се смањује. У смеши Bi₂O₃–TiO₂ механички индукована хемијска реакција доводи до образовања нанокристалне Bi₄Ti₃O₁₂ фазе током периода млевења до 1 h. Даљим млевењем ова нанокристална фаза се трансформише у аморфну фазу.

Показано је да стационарно стање (однос нанокристална/аморфна фаза) зависи од услова млевења: већи интензитет млевења индукује „нанокристализацију“ аморфне фазе, тј. таложење (издавање) кристалита величине од око 170 Å.

- [1] М. Здујић, Ч. Јовалекић, Љ. Карановић, Д. Полети, X конференција српског кристалографског друштва, Књига извода радова, Београд, 2002. с. 38.

RADIAL DISTRIBUTION FUNCTION AND THE LOCAL STRUCTURE OF SOLID SOLUTIONS OF $\text{Li}_{0.5}\text{Fe}_{2.5-x}\text{La}_x\text{O}_4$ TYPE

A. Bordaš^a, S. Rakić^a, B. Antić^b and A. Kapor^a

^a Faculty of Sciences, Trg D. Obradovića 4, 21000 Novi Sad, Serbia and Montenegro;

^b Institute of Nuclear Sciences "Vinča", P. O. Box 522, 11001 Beograd, Serbia and Montenegro

e-mail: arpi@uns.ns.ac.yu

Within the framework of the systematic analysis of the structural properties of the solid solutions of $\text{Li}_{0.5}\text{Fe}_{2.5-x}\text{La}_x\text{O}_4$ type on the basis of the powder diffractograms, a calculation was performed of the radial density function (RAD) [1] for the basic ferrite ($x=0$) and two solid solutions with lantanas ($x=0.1$ and 0.05). These materials were synthesized by the procedure for obtaining nanoparticle samples [2] and powder diffractograms recorded with $\text{CuK}\alpha$ radiation indicated a possible presence of another phase.

The aim of these calculations was to demonstrate that one can notice the difference in the local structure of the solid solution by calculating PDF (pair distribution function), which is an average representation of the atomic ordering and provides the probability of finding certain atoms at a distance r from the referent atom [3]. PDF is the representation of diffraction data, obtained in this case from nanoparticle materials, in the real space where the distance between characteristic peaks represents the separation between the pairs of atoms. By calculating PDF from the model structure and comparing it with measured values, we can obtain the information about the local structure.

Calculated values of PDF function for three studied samples showed the difference in the positions of interatomic separations in the first and higher coordination spheres with a trend of increase of this difference for higher order neighbors.

Model structure can be established rather realistically since the crystal structure of Li-ferrite is refined by the Rietveld method in the space group $P4_3,32$. The structural study of nanoparticle solid solutions showed that the decrease of symmetry occurs and an eventual presence of another phase. These uncertainties of the structure justified the attempts of the analysis of the local structure with the aid of PDF function, as well as considering its possible advantages in the preliminary phase of the study of nanoparticle materials. Observed dependence of the structure of studied materials on sample preparation procedure is another reason to apply the proposed approach to such samples.

[1] V. Petkov, *J. Appl. Cryst.* 22 (1989), 387-389.

[2] R. G. Charel, M. A. Pawlikowski, *J. Phys. Chem.*, 62 (1958), 440.

[3] H. Klug, L. Alexander, X-ray Diffraction Procedures for Polycrystalline and Amorphous Materials, Wiley, (1974), New York.

RADIJALNA FUNKCIJA RASPODELE I LOKALNA STRUKTURA ČVRSTIH RASTVORA TIPE $\text{Li}_{0.5}\text{Fe}_{2.5-x}\text{La}_x\text{O}_4$

A. Bordaš^a, S. Rakić^a, B. Antić^b and A. Kapor^a

^a Prirodno-matematički fakultet, Trg D. Obradovića 4, 21000 Novi Sad, Srbija i Crna Gora; ^b Institut za nuklearne nauke „Vinča”, p. pr. 522, 11001 Beograd, Srbija i Crna Gora
e-mail: arpi@uns.ns.ac.yu

U okviru sistematske analize strukturalnih osobina čvrstih rastvora $\text{Li}_{0.5}\text{Fe}_{2.5-x}\text{La}_x\text{O}_4$ tipa na bazi difraktograma praha urađeno je izračunavanje radijalne funkcije gustine (RAD) [1] za osnovni ferit ($x=0$) i dva čvrsta rastvora sa lantanom ($x=0,1$ i $0,05$). Ovi materijali su sintetisani postupkom za dobijanje nanočestičnih uzoraka [2] i snimljeni difraktogrami praha korišćenjem CuK α zračenja, ukazivali su na moguće prisustvo druge faze.

Cilj ovih proračuna je bio da se pokaže da je moguće uočiti razliku u lokalnoj strukturi čvrstog rastvora izračunavanjem PDF (pair distribution function), koja daje prikaz srednjeg rasporeda atoma i daje verovatnoću nalazeњa određenih atoma na rastojanju r od referentnog atoma [3]. PDF je prikaz difrakcionih podataka, u ovom slučaju dobijenih sa nanočestičnim materijala, u realnom prostoru gde rastojanje između karakterističnih pikova prikazuje razmak između parova atoma. Izračunavanjem PDF iz modelne strukture i poređenjem sa merenim vrednostima možemo dobiti informacije o lokalnoj strukturi.

Izračunate vrednosti PDF funkcije za tri ispitivana uzorka pokazala su razliku u položajima međuatomskih rastojanja u prvoj i višim koordinacionim sferama sa tendencijom povećanja te razlike za susede višeg reda.

Modelnu strukturu možemo dosta realno da postavimo jer je kristalna struktura Li-ferita utaćnjena Ritveldovom metodom u prostornoj grupi P4₃32. Strukturno ispitivanje nanočestičnih čvrstih rastvora pokazalo je da dolazi do smanjenja simetrije i eventualno postojanje druge faze. Ove nedoumice u strukturi opravdale su pokušaje analize lokalne strukture preko PDF, kao i sagledavanje eventualnih prednosti ove analize u preliminarnoj fazi ispitivanja nanočestičnih materijala. Uočena zavisnost strukture ispitivanih materijala od postupka pripreme uzorka još je jedan razlog da se predložena metoda primeni na ovakve uzorke.

- [1] V. Petkov, *J. Appl. Cryst.* 22 (1989), 387-389.
- [2] R. G. Charel, M. A. Pawlikowski, *J. Phys. Chem.*, 62 (1958), 440.
- [3] H. Klug, L. Alexander, *X-ray Diffraction Procedures for Polycrystalline and Amorphous Materials*, Wiley, (1974), New York.

NON-COVALENT INTERACTION IN CRYSTAL STRUCTURE OF E- AND Z- ISOMERS OF UNSATURATED ESTERS OF C-GLYCOSIDES

A. Kapor^a, D. Zobel^b, M. Strümpel^b, Lj. Torović^a, M. Popsavin^a

^a*Faculty of Natural Sciences and Mathematics, Trg D.Obradovića 4, 21000 Novi Sad, Serbia and Montenegro;*

^b*Institute für Kristallographie, Freie Universität, Berlin, Takustraße 6, D-14195 Berlin, Germany;
e-mail:akapor@uns.ns.ac.yu*

The unsaturated esters of E- and Z- forms were prepared like as intermediate in the process of the syntheses of target C-nucleoside. The crystal and molecular structure were studied in previous contribution [1,2]. We describe here the crystal packing and the intermolecular interaction.

It was first noticed that **b** and **c** crystallographic axes are approximately doubled in Z-isomer, which increases the unit cell volume almost four times ($V_1 = 665.98(4) \text{ \AA}^3$, $V_2=2610(1) \text{ \AA}^3$) and changes the packing symmetry and the space group ($P1 \rightarrow P2_12_12_1$).

The basic difference between E (trans) and Z (cis) isomer in the crystalline form is in the creation of a new intramolecular hydrogen bond N5-H5...O1 (3.248(1) Å) in Z-isomer leading to the formation of new seven-member quasi ring. The rest of the molecule in both isomers retains the same conformation due to strong intramolecular hydrogen contacts in which the atom O16 is a fourfold acceptor of the hydrogen from the atoms C6, C7, C15 and C22. During the crystallization process, the change of the molecular conformation induces different packing and the change of the space group of the crystal unit cell. Relatively rigid conformation of the molecule except for the side chain at C4 atom, at the first step provide the packing of free molecules parallel to each other forming chains, with similar strong intermolecular hydrogen bonds in both isomers. In the second step, molecular chains bond mutually by relatively weak hydrogen bonds of C-H...O type, which vary in two studied isomers. It was observed that the acceptors of C-H...O contacts in this case are the peripheral oxygen atoms O9 and O30.

- [1] A. Kapor, D. Zobel, M. Strümpel, Lj. Torović, M. Popsavin, Abstarcts, IX-Conference of the Serbian Crystallographic Society, (2001), 24-25
- [2] A. Kapor, D. Zobel, M. Strümpel, Lj. Torović, M. Popsavin, Abstarcts, IX-Conference of the Serbian Crystallographic Society, (2002), 19-20

NEKOVALENTNE INTERAKCIJE U KRISTALNOJ STRUKTURI E- I Z- IZOMERA NEZASIĆENOG ESTRA C-GLUKOZIDA

A. Kapor^a, D. Zobel^b, M. Strümpel^b, Lj. Torović^a, M. Popsavin^a

^a*Prirodno-matematički fakultet, Trg D.Obradovića 4, 21000 Novi Sad, Srbija i Crna Gora;*

^b*Institute für Kristallographie, Freie Universität, Berlin, Takustraße 6, D-14195 Berlin,
Nemačka;*

e-mail:akapor@uns.ns.ac.yu

Nezasićeni estar E- i Z- forme je dobijen kao međuprodukt u procesu sinteze ciljnog C-nucleozida. Rešavanje kristalne i molekulske strukture ovih izomera je analizirano u prethodnim saopštenjima [1,2]. Ovde će biti opisano pakovanje molekula u kristalnoj rešetki kao i karakteristične intermolekularne interakcije.

Prvo je uočeno da se **b** i **c** kristalografske ose približno udvostručavaju u Z-izomeru, što povećava zapreminu elementarne čelije četiri puta ($V_1 = 665,98(4) \text{ \AA}^3$, $V_2 = 2610(1) \text{ \AA}^3$) i menja simetriju pakovanja, odnosno prostornu grupu ($P1 \rightarrow P2_{12}2_1$).

Osnovna razlika između E (trans) i Z (cis) izomera u kristalnoj formi je u nastanku nove unutar-molekulske vodonične veze N5-H5...O1 ($3,248(1) \text{ \AA}$) u Z-izomeru koja dovodi do formiranja novog sedmočlanog kvazi prstena. Ostatak molekula u oba izomera zadržava istu konformaciju zahvaljujući jakim unutar molekulskim vodoničnim kontaktima u kojima je atom O16 četvorostruki akceptor vodonika od atoma C6, C7, C15 i C22. U procesu kristalizacije izmena konformacije molekula uslovjava drugačije pakovanje i promenu prostorne grupe elementarne čelije kristala. Relativno rigidna konformacija molekula, osim bočnog lanca na C4 atomu, u prvom koraku obezbeđuje pakovanje slobodnih molekula paralelno jedne drugima sa sličnim jakim međumolekulskim vodoničnim vezama u oba izomera. U drugom koraku ovako formirani lanci molekula se među sobom vezuju nešto slabijim vodoničnim vezama tipa C-H...O koje su različite u dva ispitivana izomera. Uočeno je da su akceptori C-H...O kontakata u ovom slučaju periferni atomi kiseonika O9 i O30.

- [1] A. Kapor, D.Zobel, M.Strümpel, Lj.Torović, M.Popsavin, Abstarcts, IX-Conference of the Serbian Crystallographic Society, (2001), 24-25
- [2] A. Kapor, D.Zobel, M.Strümpel, Lj.Torović, M.Popsavin, Abstarcts, IX-Conference of the Serbian Crystallographic Society, (2002), 19-20

THREE-DIMENSIONAL HYDROGEN-BONDING NETWORK IN THE CRYSTAL STRUCTURE OF PYRIDOXAL SEMICARBAZONE CHLORIDE MONOHYDRATE

G. A. Bogdanović^a, V. S. Jevtović^b, V. M. Leovac^b

^a "VINČA" Institute of Nuclear Sciences, Laboratory of Theoretical Physics and Condensed Matter Physics, 11001, Belgrade, P.O.Box 522, Serbia and Montenegro; ^b Institute of Chemistry, Faculty of Sciences, University of Novi Sad, 21000 Novi Sad, Serbia and Montenegro

e-mail: goranb@rt270.vin.bg.ac.yu

Crystal structure of pyridoxal semicarbazone chloride monohydrate, $C_9H_{13}N_4O_3 \cdot Cl \cdot H_2O$, has been determined by single crystal X-ray analysis. Data were collected on an Enraf-Nonius CAD-4 diffractometer using $MoK\alpha$ radiation ($\lambda = 0.71069 \text{ \AA}$) and $\omega/2\theta$ scans in 2.13 to 26.96° θ range. Cell constants and an orientation matrix for data collection, obtained from 24 centered reflections in the θ range 12.30 - 15.61° corresponded to a triclinic system. The data were corrected for Lorentz and polarization effects. The structure was solved by direct methods and difference-Fourier methods, and refined by full matrix least-squares using SHELXL97 program to a final $R = 0.0635$ for 1851 reflections with $I > 2\sigma(I)$. Water molecule and hydroxyl H-atom positions were determined by the HYDROGEN program [1] and were refined using a riding model with a fixed O-H bond length of 0.85 \AA ; isotropic displacement parameters of hydrogen atoms were set equal to 1.5 times the equivalent isotropic displacement parameter of the parent atoms.

All non-hydrogen atoms from the cation of pyridoxal semicarbazone (PxSC) are approximately coplanar. The presence of chloride anions and water molecules gives rise to an extensive hydrogen-bonded system. The crystal packing consists of parallel layers which are oriented in the direction of PxSC plane. These layers are interconnected by hydrogen bonds through the Cl anions and water molecules. Stereochemistry of presented crystal structure is compared to previously published structures of pyridoxal thiosemicarbazone trihydrate [2] and pyridoxal thiosemicarbazone chloride [3].

Crystallographic data: $C_9H_{15}ClN_4O_4$, triclinic, space group $P\bar{1}$, $a = 7.163(2)$, $b = 9.1320(10)$, $c = 10.310(2) \text{ \AA}$, $\alpha = 70.750(10)$, $\beta = 77.49(2)$, $\gamma = 81.462(10)^\circ$, $V = 619.4(2) \text{ \AA}^3$, $Z = 2$, $D_c = 1.494 \text{ Mg/m}^3$, $\mu(MoK\alpha) = 0.323 \text{ mm}^{-1}$, $MoK\alpha$ radiation, $\lambda = 0.71069 \text{ \AA}$.

- [1] M. Nardelli, *J. Appl. Cryst.*, 32 (1999) 563.
- [2] U. Abram, K. Ortner, R. Gust, K. Sommer, *J. Chem. Soc., Dalton Trans.*, (2000) 735.
- [3] M. F. Belicchi, G. F. Gasparri, E. Leporati, C. Pelizzi, P. Tarasconi, G. Tosi, *J. Chem. Soc., Dalton Trans.*, (1986) 2455.

TRODIMENZIONALNA MREŽA VODONIČNIH VEZA U KRISTALNOJ STRUKTURI PIRIDOKSAL-SEMIKARBAZON-HLORID-MONOHIRDRATA

G. A. Bogdanović^a, V. S. Jevtović^b, V. M. Leovac^b

^a*Institut za nuklearne nauke "VINČA", Laboratorija za teorijsku fiziku i fiziku kondenzovane materije, 11001, Beograd, p. p. 522, Srbija i Crna Gora;* ^b*Prirodno-matematički fakultet, Univerzitet u Novom Sadu, Trg Dositeja Obradovića 3, 21000 Novi Sad, Srbija i Crna Gora*
e-mail: goranb@rt270.vin.bg.ac.yu

Kristalna struktura piridoksal-semikarbazone-hlorid-monohidrata, $C_9H_{13}N_4O_3 \cdot Cl \cdot H_2O$, određena je rendgenskom struktturnom analizom. Eksperimentalni podaci su prikupljeni na difraktometru uz korišćenje monohromatskog $MoK\alpha$ zračenja ($\lambda = 0,71069 \text{ \AA}$) i $\omega/2\theta$ skeniranja u opsegu θ ugla od $2,13$ do $26,96^\circ$. Dimenzije jedinične celije i orijentaciona matrica dobijeni su korišćenjem 24 centriranih refleksija koje su bile u opsegu θ ugla od $12,30$ - $15,61^\circ$ i odgovarale su trikliničnom sistemu. Podaci su korigovani na Lorensov i polarizacioni faktor. Struktura je rešena direktnim metodama i differentnom Furijeovom sintezom a utačnjena je metodom najmanjih kvadrata pune matrice korišćenjem programa SHELLXL97 do finalnog $R = 0,0635$ za 1851 refleksiju koje zadovoljavaju uslov $I > 2\sigma(I)$. Atomi vodonika iz molekula vode i hidroksilnih grupa određeni su korišćenjem programa HYDROGEN [1], a utačnjeni su sa fiksiranim dužinama O-H veza u vrednosti od $0,85 \text{ \AA}$. Izotropni temperaturni faktori ovih atoma vodonika fiksirani su na vrednost $1,5$ puta veću od vrednosti ekvivalentnih izotropnih parametara nevodonikovih atoma za koje su vezani.

Svi nevodonikovi atomi iz katjona piridoksal-semikarbazone ($PxSC$) su približno koplanarni. Prisustvo anjona hlora i molekula vode omogućilo je građenje sistema vodoničnih veza. Kristalno pakovanje se sastoji od paralelnih slojeva orijentisanih u pravcu pružanja $PxSC$ -ravnji. Ovi slojevi su međusobno povezani građenjem vodoničnih veza preko anjona hlora i molekula vode. Stereohemija prezentovane kristalne strukture upoređena je sa prethodno publikovanim strukturama piridoksal-tiosemikarbazone-trihidrata [2] i piridoksal-tiosemikarbazone-hlorida [3].

Kristalografski podaci: $C_9H_{15}ClN_4O_4$, triklinični sistem, prostorna grupa $P\bar{1}$, $a = 7,163(2)$, $b = 9,1320(10)$, $c = 10,310(2) \text{ \AA}$, $\alpha = 70,750(10)$, $\beta = 77,49(2)$, $\gamma = 81,462(10)^\circ$, $V = 619,4(2) \text{ \AA}^3$, $Z = 2$, $D_c = 1,494 \text{ Mg/m}^3$, $\mu(MoK\alpha) = 0,323 \text{ mm}^{-1}$, $MoK\alpha$ zračenje, $\lambda = 0,71069 \text{ \AA}$.

- [1] M. Nardelli, *J. Appl. Cryst.*, 32 (1999) 563.
- [2] U. Abram, K. Ortner, R. Gust, K. Sommer, *J. Chem. Soc., Dalton Trans.*, (2000) 735.
- [3] M. F. Belicchi, G. F. Gasparri, E. Leporati, C. Pelizzetti, P. Tarasconi, G. Tosi, *J. Chem. Soc., Dalton Trans.*, (1986) 2455.

SYNTHESIS AND CRYSTAL STRUCTURE OF 3,16,17-TRIHYDROXY-16,17-SECOESTRA-1,3,5(10)-TRIENE


D. Lazar^a, S. Stanković^a, O. Klisurić^a, S. Jovanović-Šanta^a, J. Petrović^a

^a*Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia and Montenegro
e-mail: dlazar@uns.ns.ac.yu*

In order to study estrogenic vs. antiestrogenic activity, a new compound, 3,16,17-trihydroxy-16,17-secoestra-1,3,5(10)-triene, was synthesized in several synthetic steps starting from estrone.

The compound crystallizes in the orthorhombic system, space group P2₁2₁2₁, with the unit cell parameters $a = 10.195(5)$, $b = 6.675(5)$, $c = 23.145(5)$ Å, $Z = 4$, $D_x = 1.225$ M gm⁻³, $\mu = 0.08$ mm⁻¹, MoK_α radiation.

The crystal structure was solved by direct methods on the basis of 2515 independent reflections using SHELXS-97 [1] and refined by SHELXL-97 [1]. The final R factor is 0.0216 for 1027 reflections with $I > 2\sigma(I)$.



Perspective view of the molecule

- [1] G. M. Sheldrick, (1997) SHELX-97, *Program for the solution and refinement of crystal structures*, University of Göttingen, Germany.

SINTEZA I KRISTALNA STRUKTURA 3,16,17-TRIHIDROKSI-16,17-SEKOESTRA-1,3,5,(10)-TRIENA


D. Lazar^a, S. Stanković^a, O. Klisurić^a, S. Jovanović-Šanta^a, J. Petrović^a

^aPrirodno-matematički fakultet, Univerzitet u Novom Sadu, Trg Dositeja Obradovića 4,
21000 Novi Sad, Srbija i Crna Gora
e-mail: dlazar@uns.ns.ac.yu

U cilju ispitivanja estrogene odnosno antiestrogene aktivnosti, polazeći od estrona, sintetizovano je u više sintetskih faza novo jedinjenje:
3,16,17-trihidroksi-16,17seko-estra-1,3,5(10)-trien.

Jedinjenje kristališe u ortorombičnom kristalografskom sistemu, prostorna grupa P2₁2₁2₁, sa parametrima elementarne celiye a = 10,195(5), b = 6,675(5), c = 23,145(5) Å, Z = 4, D_x = 1,225 Mg m⁻³, μ = 0,08 mm⁻¹, MoK_α zračenje.

Struktura kristala rešena je primenom direktnih metoda na bazi 2515 nezavisnih refleksa pomoću SHELXS-97 [1] i utaćnjena pomoću SHELXL-97 [1]. Konačna vrednost R-faktora je 0,0216 za 1027 refleksa sa $I > 2\sigma(I)$.



Trodimenzionalni prikaz molekula


[1] G. M. Sheldrick, (1997) SHELX-97, *Program for the solution and refinement of crystal structures*, University of Göttingen, Germany.

CRYSTAL STRUCTURE AND THE INFLUENCE OF NEW 3,17-SUBSTITUED D-SECO-ESTRONE DERIVATIVES ONTO THE PHYSICAL CHARACTERISTICS OF THE BINARY MIXTURES OF THE CHOLESTERIC LIQUID CRYSTALS

M. Garić^a, D. Ž. Obadović^a, S. Jovanović-Šanta^b, D. Lazar^a and A. Vajda^c

^a*Department of Physics, Faculty of Sciences, Trg D. Obradovića 4, Novi Sad;* ^b*Department of Chemistry, Faculty of Sciences, Trg D. Obradovića 3, Novi Sad, Serbia and Montenegro;*
^cResearch Institute for Solid State Physics and Optics of the Hungarian Academy of Sciences, H-1525 Budapest, P.O.Box 49, Hungaria
e-mail: gara@im.ns.ac.yu

The crystal structure and influence of the new chiral additives, potential antiestrogenes, onto the physical characteristics of the binary mixtures of cholesteric liquid crystals were studied. The tested D-seco-estratriene-derivatives **1-3** (3-benzyloxy-17-oxo-16,17-secoestra-1,3,5(10)-triene-16-nitrile, **1**; 3-benzyloxy-17-hydroxy-16,17-secoestra-1,3,5(10)-triene-16-nitrile, **2** and 3-hydroxy-17-*p*-toluenesulphonyloxy-16,17-secoestra-1,3,5(10)-triene-16-nitrile, **3**) were synthesized in several synthetic steps, starting from estrone.



The phase transitions of the mixture of cholesteryl laurate and cholesteryl enantate with the added chiral additives **1-3** (45%-45%-10%; **Mix1**, **Mix2** and **Mix3**, respectively) were examined. A considerable shift of the phase transition temperature I→Ch, as well as the temperature of the SmA phase formation was observed. X-ray diffraction data enabled the determination of the crystal and molecular structure of the compound **3** and its molecular length (*l*) and width (*d**), as well as the molecular parameters of the mixtures **Mix1**, **Mix2** and **Mix3**: the thickness of smectic *i.e.* long spacing distance of cholesteric layers (*d*) and average distance between the long axes of neighbouring molecules (*D*). The most probably conformation of the compounds **1**, **2** were determined by the molecular-mechanic calculations on the basis of solved crystal structure. The influence of these chiral additives onto the pitch of the cholesteric helix of the studied binary mixtures of the cholesteric liquid crystals was also investigated.


KRISTALNA STRUKTURA I UTICAJ NOVIH 3,17-SUPSTITUISANIH D-SEKO-ESTRONSKIH DERIVATA NA FIZIČKE KARAKTERISTIKE BINARNIH SMEŠA HOLESTERIČKIH TEČNIH KRISTALA

M. Garić^a, D. Ž. Obadović^a, S. Jovanović-Šanta^b, D. Lazar^a i A. Vajda^c

^aDepartman za fiziku, Prirodno-matematički fakultet, Trg D. Obradovića 4, Novi Sad;

^bDepartman za hemiju, Prirodno-matematički fakultet, Trg D. Obradovića 3, Novi Sad, Srbija i Crna Gora; ^cIstraživački institut za fiziku čvrstog stanja i optiku Mađarske Akademije nauka, H-1525 Budimpešta, p. pr. 49, Mađarska
e-mail: gara@im.ns.ac.yu

Određena je kristalna struktura i ispitivan uticaj novih hiralnih aditiva, potencijalnih antiestrogena, na fizičke karakteristike binarnih smeša holesteričkih tečnih kristala. Ispitivani D-seko-estratriensi derivati **1-3** (3-benziloksi-17-okso-16,17-sekoestra-1,3,5(10)-trien-16-nitril, **1**; 3-benziloksi-17-hidroksi-16,17-sekoestra-1,3,5(10)-trien-16-nitril, **2** i 3-hidroksi-17-*p*-toluensulfoniloksi-16,17-sekoestra-1,3,5(10)-trien-16-nitril, **3**) sintetisani su u nekoliko sintetskih faza, polazeći od estrona.



Ispitivani su fazni prelazi smeša holesteril laurata i holesteril enantata prilikom dodavanja hiralnih aditiva **1-3** (45%-45%-10%; **Mix1**, **Mix2** and **Mix3**, respektivno). Primećen je značajan pomak temperature faznog prelaza I→Ch, kao i temperature formiranja SmA faze. Podaci dobijeni difrakcijom x-zraka omogućili su određivanje kristalne i molekulske strukture, tj. dužine (l) i širine (d*) molekula jedinjenja **3**, kao i molekulske parametara smeša **Mix1**, **Mix2** i **Mix3**: debljine smektičkih slojeva, odnosno uređenja na daljinu holesteričkih slojeva (d) i srednje vrednosti rastojanja između dugih osa susednih molekula (D). Najverovatnija konformacija jedinjenja **1** i **2** je određena pomoću molekularno-mehaničkih proračuna na bazi kristalne strukture jedinjenja **3**. Takođe je ispitivan uticaj hiralnih aditiva na promenu koraka holesteričke spirale binarnih smeša holesteričkih tečnih kristala.

SYNTHESIS AND CRYSTAL STRUCTURE INVESTIGATIONS OF TWO ANDROSTENE DERIVATIVES

O. Klisurić^a, S. Stanković^a, D. Lazar^a, D. Zobel^b, K. Penov-Gašić^a, S. Stajanović^a


^aFaculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia and Montenegro, ^bInstitute für Chemie/Kristallographie, Freie Universität, Berlin, Takustrasse 6, D-14195 Berlin, Germany
e-mail: olivia@uns.ns.ac.yu

17 α -methyl-17-oxa-D-homo-4-androstene-3,16-dione (**1**) and 17 α -methyl-17-oxa-D-homo-1,4,6-androstatrien-3,16-dione (**2**) were synthesized from 3 β -hydroxy-16-oximino-5-androstene-17-one in several synthetic steps.


The compound **1** crystallizes in the triclinic system, space group P1, with the unit cell parameters $a=8.040(2)$, $b=8.160(2)$, $c=13.658(7)$ Å; $\alpha=77.03(3)$, $\beta=88.95(3)$, $\gamma=89.22(2)$ °, $V=854.95(8)$ Å³, $Z=2$, $D_x=1.229$ M gm⁻³, $\mu=0.08$ mm⁻¹, MoK α radiation.

The compound **2** crystallizes in the orthorhombic system, space group P2₁2₁2₁, with the unit cell parameters $a=7.225(2)$, $b=8.887(2)$, $c=25.358(8)$ Å, $V=1628.2(2)$ Å³, $Z=4$, $D_x=1.270$ M gm⁻³, $\mu=0.08$ mm⁻¹, MoK α radiation.

Both structures were solved by direct methods using SHELXS-97 [1], and refined by SHELXL-97 [1]. The final R factors for compounds **1** and **2** are 0.0486 (for 3683 reflections with I>2σ(I)) and 0.0427 (for 1547 reflections with I>2σ(I)), respectively.



Perspective view of the molecule **1**



Perspective view of the molecule **2**

[1] G. M. Sheldrick, SHELX-97, *Program for the solution and refinement of crystal structures*, University of Goettingen, Germany (1997).

SINTEZA, STRUKTURNA I KRISTALOGRAFSKA ISTRAŽIVANJA DVA DERIVATA ANDROSTENA

O. Klisurić^a, S. Stanković^a, D. Lazar^a, D. Zobel^b, K. Penov-Gašić^a, S. Stajanović^a


^a Prirodno-matematički fakultet, Univerzitet u Novom Sadu, Trg Dositeja Obradovića 4, 21000 Novi Sad, Srbija i Crna Gora, ^b Institute für Kristallographie, Freie Universität, Berlin, Takustrasse 6, D-14195 Berlin, Nemačka
e-mail: olivia@uns.ns.ac.yu

17 α -metil-17-oksa-D-homo-4-androsten-3,16-dion (**1**) i 17 α -metil-17-oksa-D-homo-1,4,6-androstriien-3,16-dione (**2**) su sintetizovani polazeći od 3 β -hidroksi-16-oksimino-5-androsten-17-ona u nekoliko sintetskih faza.

Jedinjenje **1** kristališe u triklinskom kristalografskom sistemu, prostorna grupa P1, sa parametrima elementarne celije $a=8,040(2)$, $b=8,160(2)$, $c=13,658(7)$ Å; $\alpha=77,03(3)$, $\beta=88,95(3)$, $\gamma=89,22(2)$ °, $V=854,95(8)$ Å³, $Z=2$, $D_x=1,229$ M g m⁻³, $\mu=0,08$ mm⁻¹, MoKα zračenje.

Jedinjenje **2** kristališe u ortorombičnom kristalografskom sistemu, prostorna grupа P2₁2₁2₁, sa parametrima elementarne celije $a=7,225(2)$, $b=8,887(2)$, $c=25,358(8)$ Å, $V=1628,2(2)$ Å³, $Z=4$, $D_x=1,270$ Mg m⁻³, $\mu=0,08$ mm⁻¹, MoKα zračenje.

Obe strukture su rešene direktnim metodama primenom SHELXS-97 [1] i utaćnjene pomoću SHELXL-97 [1]. Konačni R faktori za jedinjenja **1** and **2** su 0,0486 (za 3683 refleksa sa $I>2\sigma(I)$) i 0,0427 (za 1547 refleksa sa $I>2\sigma(I)$), respektivno.



[1] G. M. Sheldrick, *SHELX-97, Program for the solution and refinement of crystal structures*, University of Goettingen, Germany (1997).

SYNTHESIS AND CRYSTAL STRUCTURE OF (17R)-3,17-DIHYDROXY-16-CYANO-17-METHYL-16,17-SECOESTRA-1,3,5(10)-TRIENE (1) AND 3-BENZYLOXY-17A-METHYL-17-OXA-D-HOMOEstra-1,3,5-TRIEN-16-ONE (2)

O. Klisurić^a, S. Stanković^a, D. Lazar^a, D. Zobel^b, A. Wagner^b, K. Penov-Gašić^a, S. Stajanović^a



^aFaculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia and Montenegro, ^bInstitute für Chemie/Kristallographie, Freie Universität, Berlin, Takustrasse 6, D-14195 Berlin, Germany
e-mail: olivia@uns.ns.ac.yu

In order to investigate anti-hormonal activity we were synthesized (17R)-3,17-dihydroxy-16-cyano-17-methyl-16,17-secoestra-1,3,5(10)-triene (**1**) starting from estrone benzyl ether in five synthetic steps. In one of the synthetic steps we were obtained 3-benzyloxy-17a-methyl-17-oxa-D-homoestra-1,3,5-trien-16-one (**2**) as by product.

The compound **1** crystallizes in the orthorhombic system, space group P2₁2₁2₁, with the unit cell parameters $a=8.218(2)$, $b=9.567(2)$, $c=21.383(7)$ Å; $V=1681.0(8)$ Å³, $Z=4$, $D_x=1.183$ Mg m⁻³, $\mu=0.08$ mm⁻¹, MoKα radiation.

The compound **2** crystallizes in the triclinic system, space group P1, with the unit cell parameters $a=6.852(2)$, $b=8.744(2)$, $c=9.740(2)$ Å; $\alpha=91.903(3)$, $\beta=101.881(3)$, $\gamma=112.525(3)$ °, $V=523.38(8)$ Å³, $Z=1$, $D_x=1.239$ Mg m⁻³, $\mu=0.08$ mm⁻¹, MoKα radiation.

Both structures were solved by direct methods using SHELXS-97 [1], and refined by SHELXL-97 [1]. The final R factors for compounds **1** and **2** are 0.0349 (for 5199 reflections with $I>2\sigma(I)$) and 0.0354 (for 6305 reflections with $I>2\sigma(I)$), respectively.



[1] G. M. Sheldrick, SHELXL-97, *Program for the refinement of crystal structures*, University of Goettingen, Germany (1997).

SINTEZA I KRISTALNA STRUKTURA (17R)-3,17-DIHIDROKSI-16-CIJANO-17-METIL-16,17-SEKOESTRA-1,3,5(10)-TRIENA (1) I 3-BENZILOKSI-17A-METIL-17-OKSA-D-HOMO-ESTRA-1,3,5-TRIEN-16-ONA (2)

O. Klisurić^a, S. Stanković^a, D. Lazar^a, D. Zobel^b, A. Wagner^b, K. Penov-Gašić^a, S. Stajanović^a


^aPrirodno-matematički fakultet, Univerzitet u Novom Sadu, Trg Dositeja Obradovića 4, 21000 Novi Sad, Srbija and Crna Gora, ^bInstitute für Chemie/Kristallographie, Freie Universität, Berlin, Takustrasse 6, D-14195 Berlin, Nemačka
e-mail: olivia@uns.ns.ac.yu

U cilju ispitivanja antihormonske aktivnosti sintetizovan je (17R)-3,17-dihidroksi-16-cijano-17-metil-16,17-sekoestra-1,3,5(10)-trien (**1**) iz benziletra estrona u pet sintetskih faza. U jednoj od sintetskih faza kao sporedan proizvod reakcije dobijen je 3-benziloksi-17a-metil-17-oksa-D-homoestra-1,3,5-trien-16-on (**2**).

Jedinjenje (**1**) kristališe u ortorombičnom kristalografskom sistemu, prostorna grupa P2₁2₁2₁, sa parametrima elementarne čelije $a=8,218(2)$, $b=9,567(2)$, $c=21,383(7)$ Å; $V=1681,0(8)$ Å³, $Z=4$, $D_x=1,183$ M gm⁻³, $\mu=0,08$ mm⁻¹, MoKα zračenje

Jedinjenje (**2**) kristališe u triklinском kristalografskom sistemu, prostorna grupa P1, sa parametrima elementarne čelije $a=6,852(2)$, $b=8,744(2)$, $c=9,740(2)$ Å; $\alpha=91,903(3)$, $\beta=101,881(3)$, $\gamma=112,525(3)$ °, $V=523,38(8)$ Å³, $Z=1$, $D_x=1,239$ M gm⁻³, $\mu=0,08$ mm⁻¹, MoKα zračenje.

Obe strukture su rešene direktnim metodama primenom SHELXS-97 [1] i utačnjene pomoću SHELXL-97 [1]. Konačni R-faktori za jedinjenja **1** and **2** su 0,0349 (za 5199 reflections sa $I>2\sigma(I)$) i 0,0354 (za 6305 reflections sa $I>2\sigma(I)$), respektivno.



[1] G. M. Sheldrick, SHELXL-97, *Program for the refinement of crystal structures*, University of Goettingen, Germany (1997).